

Fax Referral To: 1-855-297-1270

Address: 280 Avenida Jesus T. Pinero Ste B Rio Piedras, PR 00927

Phone: 1-888-280-1190

NCPDP: 4026325

Six Simple Steps to Submitting a Referral **PATIENT INFORMATION** (Complete or include demographic sheet) Patient Name: _____ DOB: __ Address: ____City, State, ZIP Code: ___ Preferred Contact Methods: Phone (to primary # provided below) Text (to cell # provided below) Email (to email provided below) Note: Carrier charges may apply. By providing the phone number(s) and email address above, you are consenting to receive automated calls, emails and/or text messages from CVS Specialty® about your prescription(s), account, and health care. Standard data rates apply. Message frequency varies. If unable to contact via text or email, Specialty Pharmacy will attempt to contact by phone. Primary Phone: ______ Alternate Phone: _____ Email: _____ Last Four of SSN: _____ Primary Language: _____ Parent/Caregiver/Legal Guardian Name (Last, First): _______Relationship to patient: _____ 2 PRESCRIBER INFORMATION Prescriber's Name: ______ DEA #: _____ Group or Hospital: _____ Prescriber's Name: ___ _____ State License #: ______ Address: ______ City, State, ZIP Code: ______ Phone: _____ Contact Person: _____ Contact's Phone: _____ INSURANCE INFORMATION Please fax copy of prescription and insurance cards with this form, if available (front and back) 4 DIAGNOSIS AND CLINICAL INFORMATION Needs by Date: Ship to: Patient Office Other: Diagnosis (ICD-10): Date of Diagnosis / / K50.00 Crohn's Disease of Small Intestine Without Complications K51.90 Ulcerative colitis, unspecified, without complications L40.50 Arthropathic Psoriasis, Unspecified L40.54 Juvenile Psoriatic Arthritis (JPsA) M06.9 Rheumatoid Arthritis. Unspecified M08.00 Juvenile Idiopathic Arthritis (JIA) M32.1 Systemic lupus erythematosus (SLE) M32.14 Glomerular disease in systemic lupus erythematosus M45.9 Ankylosing Spondylitis of Unspecified Sites in Spine M45.A0 Non-Radiographic Axial Spondylarthritis (nr-axSpA) Other Code: ______Description: _____ Patient Clinical Information:

Allergies: _____ NKDA Weight: ____ kg _ lb Height: ___

Treatment status: New to therapy Continuation of therapy; Date of last treatment __/__/__ ☐ NKDA Weight: ____ ☐ kg ☐ lb Height: ____ ☐ cm ☐ in TB Test Date __/__/ Positive Negative Hepatitis status: _____ Prior therapy, treatment dates, and reason(s) for discontinuation: _____________ **Nursing and Administration:** First dose administration of monoclonal antibodies (mABs) should be administered in a controlled setting (may vary depending upon medication specific policy). For Remicade/Remicade Biosimilars, the first dose must be administered in a controlled setting. Specialty pharmacy to coordinate home health Infusion nurse visit as necessary?

Yes
No Site of Care: Home Infusion* Coram Ambulatory Infusion Suite (AIS)* Prescriber's Office** Other Infusion Clinic *Home Infusion/Coram AIS: Diluents, Flushes, Supplies, Nursing Services for drug administration/therapy teach train. **Prescriber's Office/Other Infusion Clinic: Drug only for facility administration

<u> </u>		Please Complete Patient and F			
Patient Name: _		Patient DOB:	Patient Phone:		
			escriber Phone:		
Patient Clinical		_			
Allergies:		NKDA W	eight: 🗌 kg 🗌 lb Height: 🔲 c	m 🗌 in	
		Continuation of therapy; D			
			s status:		
Prior therapy, tre	atment dates, and re	eason(s) for discontinuation:			
	ON INFORMATION	*			
MEDICATION	STRENGTH	DOSE &	DIRECTIONS	QUANTITY/REFILLS	
_	☐ 80 mg/4 mL	☐ Induction Dose: Infuse 4 mg/kg eve	erv 4 weeks	Quantity:	
Actemra Actemra	200 mg/10 mL	Maintenance Dose: Infuse 8 mg/kg		Quantity: Refills:	
	☐ 400 mg/20 mL		over, i weeke	Keillis.	
		Ankylosing Spondylitis Induction Do	ose: Infuse IV at 5 mg/kg		
		(Dose =mg) at weeks 0, 2, 6 and e	very 6 weeks thereafter		
		Ankylosing Spondylitis Maintenance	<u>e Dose</u> : Infuse IV at 5 mg/kg		
		(Dose =mg) every 6 weeks			
			c ≥ 6 years old) <u>Induction Dose</u> : Infuse IV at		
		5 mg/kg (Dose =mg) at weeks	-		
		Crohn's Disease (Adult) Maintenand (Dose = mg) every 8 weeks	ce Dose: Infuse IV at 5-10 mg/kg		
		Crohn's Disease (Pediatric ≥ 6 years	s old) Maintenance Dose:		
		Infuse IV at 5 mg/kg (Dose =n	·		
		Plaque Psoriasis & Psoriatic Arthritis		Quantity:	
Avsola	100 mg vial	(Dose =mg) at weeks 0, 2, 6 ar		# of 100 mg vial(s)	
		Plaque Psoriasis & Psoriatic Arthritis		Refills:	
		Infuse IV at 5 mg/kg (Dose =n			
		Rheumatoid Arthritis Induction Dose	e: Infuse IV at 3 mg/kg		
		(Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis <u>Maintenance Dose</u> : Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter			
			ic \geq 6 years old) Maintenance Dose: Infuse IV	ISA IV	
		at 5 mg/kg (Dose =mg) every			
	☐ 120 mg 5 mL	at a ring, kg (Base =ring, avar)	o weeks		
□ p t t.	vial	Induction Dose: 10 mg/kg IV (Dose =mg) at 2-week intervals for the first 3		Quantity: vials	
Benlysta	☐ 400 mg 20 mL			Refills:	
	vial				
		Loading Dose:			
	125 mg/5 mL vial			Quantity:	
		Infuse 6 mg/kg (Dose = mg) at Week 0		Refills: 0	
☐ Cosentyx		Maintenance Dose: ☐ Infuse 1.75 mg/kg (Dose = mg) every 4 weeks (max. maintenance dose: 300 mg per infusion) ☐ Induction Dose: 300 mg infused IV over 30 minutes at 0, 2 and 6 weeks, then			
				Quantity:	
				Refills:	
	000			0	
□ Entrado	300 mg in a single	every 8 weeks thereafter Maintenance Dose: 300 mg infused IV over 30 minutes at 0, 2 and 6 weeks, then		Quantity:	
☐ Entyvio	dose vial in individual carton			Refills:	
	Strength:		<u> </u>	Quantity:	
Other	Suerigui.	Dose:		Refills:	
PRESCRIP	ER SIGNATI IPF	REQUIRED (STAMP SIGNAT	URE NOT ALI OWED)		
		•	,		
"Dispense As Writte DAW / May Not Sub	•	essary / Do Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible		
•	ignature:	Date:	Prescriber's Signature:	Date:	
	.gw.w. V				

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

Potiont Name:		Please Complete Patient and Prescriber Information	
	 ie:	Patient DOB: Patient Phone: Prescriber Phone:	
Prescriber Nam Patient Clinica		Prescriber Phone:	
Allergies: Treatment statu TB Test Date Prior therapy, tre	s: New to therap	reason(s) for discontinuation:	
	ON INFORMATION		
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS
☐ Inflectra	100 mg vial	Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: # of 100 mg vial(s) Refills:
Omvoh	300 mg/15 mL single dose vial	Induction Dose ☐ Week 0: Infuse 300 mg via IV infusion over at least 30 minutes ☐ Week 4: Infuse 300 mg via IV infusion over at least 30 minutes ☐ Week 8: Infuse 300 mg via IV infusion over at least 30 minutes	Quantity: Refills: 0 1 Vial 2 Vials 3 Vials
☐ Orencia ☐ Remicade ☐ Renflexis	250 mg vial	☐ Infuse mg at weeks 0, 2 and 4, then every 4 weeks thereafter ☐ Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter ☐ Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks ☐ Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks ☐ Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks ☐ Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks ☐ Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) ☐ Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter ☐ Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: Refills: Quantity: # of 100 mg vial(s) Refills:
Other	Strength:	□ Dose:	Quantity: Refills:
"Dispense As Writt DAW / May Not Su Prescriber's S	en" / Brand Medically Ned bstitute ignature:	Date:ATTN: New York and lowa providers, ples	

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

	_	Please Complete Patient and F	Prescriber Information	
Patient Name: _			Patient Phone:	
Prescriber Name			escriber Phone:	
Patient Clinical	Information:			
Allergies:		NKDA W	'eight: 🗌 kg 🗌 lb Height:	☐ cm ☐ in
		Continuation of therapy; D		
			s status:	
	ON INFORMATIO			
MEDICATION	STRENGTH	DOSE &	DIRECTIONS	QUANTITY/REFILLS
☐ Riabni ☐ Rituxan ☐ Ruxience	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separ	rated by 2 weeks	Quantity: Refills:
Saphnelo	300 mg/2 mL (150 mg/mL)	300 mg IV over a 30-minute period	d, every 4 weeks	Quantity: vials Refills:
☐ Simponi ARIA	50 mg/4 mL single dose vial	Adult RA, PsA, AS Induction Dose Week 0: Infuse 2 mg/kg IV (Dose= Week 4: Infuse 2 mg/kg IV (Dose=	=	Quantity: vials Refills:0 Quantity: vials Refills:0
		Adult RA, PsA, AS Maintenance Dos Infuse 2 mg/kg IV (Dose=mg		Quantity: vials Refills:
		Peds JIA or PsA (≥2 years old) Induc Week 0: Infuse 80 mg/m² IV (Dose Week 4: Infuse 80 mg/m² IV (Dose	=mg) over 30 minutes	Quantity: vials Refills:O Quantity: vials Refills:O
		Peds JIA or PsA (>2 years old) Maint ☐ Infuse 80 mg/m² IV (Dose=m		Quantity: vials Refills:
Skyrizi	600 mg/10 mL (60 mg/mL) single dose vial	Induction Dose: Week 0: Infuse 600 mg IV over at I Week 4: Infuse 600 mg IV over at I Week 8: Infuse 600 mg IV over at I	east one hour	Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
Stelara	130 mg/26 mL (5 mg/mL) IV single-dose vial	Single IV Induction Dose: 55 kg or less 260 mg at week 0: # of vials to be used 2 more than 55 kg to 85 kg 390 mg at week 0: # of vials to be used 3 more than 85 kg 520 mg at week 0: # of vials to be used 4		Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
☐ Truxima	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separated by 2 weeks		Quantity: Refills:
Other	Strength:	Dose:		Quantity: Refills:
	_	REQUIRED (STAMP SIGNAT	LIDE NOT ALLOWED	
"Dispense As Writte DAW / May Not Sub Prescriber's Si	en" / Brand Medically Necessostitute ignature:	eary / Do Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature: ATTN: New York and Iowa providers,	Date:

L. The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

Autoimmune IV Enrollment Form Nursing Orders

ationt Nomes:			Prescriber Information	
atient Name:		Patient DOB:	Patient Phone: Prescriber Phone:	
escriber Name:	•	P	rescriber Phone:	
atient Clinical Informat			Veight: 🗌 kg 🔲 lb Height:	□om□in
eatment status: \(\int \text{New} \)	to therapy	Continuation of therapy: I	Veight:	
Test Date//	Positive N	egative Henati	tis status:	
) for discontinuation:		
PRESCRIPTION INFO			VILL ONLY BE SENT FOR INFUSIONS DON	IE AT HOME/CODAM AIS**
MEDICATION/SUPPLIES			ENGTH/ DIRECTIONS	QUANTITY/REFILLS
ILDICATION/ SOPPLIES	ROOTE		on drug admin days – SASH or PRN to	QUANTIT I/REFIELS
Catheter: PIV PORT CVC/PICC	IV	maintain IV access and pate PIV: NS 5 mL (Heparin 10 uni CVC/PICC: NS 10 mL & H 3-5 mL.	ncy its/mL 3-5 mL if multiple days) Ieparin 10 units/mL or 100 units/mL access PORT w/ huber needle	Quantity: Refills:
		THE TO THE ATTORACTION	oo ama me o ome.	Hydration max infusion
Hydration: □ NS □ D5W	IV	Pre: 500 mL 1000 mL Concurrent: 500 mL 1000 mL Post: 500 mL 1000 mL	000 mL Other:	rate mL/hr (Adult max rate 250 mL/hr unless otherwise indicated)
☐Epinephrine *nursing requires**	□ IM □ SC	1:1000, 0.3mg/0.3 mL (gr 1:1000, 0.15mg/0.3 mL (1 1:1000, 0.1 mg/kg, Max 0 Mild-Moderate Reactions. M for severe allergic reaction, a	5-30 kg/33-66 lbs) .3mg (under 15kg) 1ay repeat in 3-5 minutes as needed	Quantity: Refills:
Diphenhydramine Oral	РО	Premedication: 12.5 mg/kg (0-30 kg) 25 mg 50 mg (Over 30 kg)		Quantity: Refills:
Diphenhydramine one one one one one one one one one o	Slow IV	1 mg/kg (under 15 kg) 12.5 mg-50 mg (15-30 kg) 25 mg-50 mg (Over 30 kg	g) nay repeat in 3-5 minutes as needed ay)	Quantity: Refills:
☐ Flush Orders:	Peripheral Access Central Venous Access	□ 10 mL NS post flush □ 50 mL NS post flush to clear medication from tubing (recommended if no post-hydration) □ Other:		Send quantity sufficient for medication days supply
Additional				
ledication:				
Patient is interested in patient supp		STAMP SIGNATURE NOT ALLOWED JIRED (STAMP SIGNAT	, ,,	provided as needed for administration
"Dispense As Written" / Brand M DAW / May Not Substitute	edically Necessary / D	o Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:
Prescriber's Signature:				

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA

request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Specialty and/or one of its

affiliates.