

Fax Referral To: 1-877-232-5455 Address: 500 Ala Moana Blvd., Bldg 1 Honolulu, HI 96813

Six Simple Steps to Submitting a Referral PATIENT INFORMATION (Complete or include demographic sheet) Patient Name: _____ DOB: __ ____City, State, ZIP Code: ___ Address: Preferred Contact Methods: Phone (to primary # provided below) Text (to cell # provided below) Email (to email provided below) Note: Carrier charges may apply. By providing the phone number(s) and email address above, you are consenting to receive automated calls, emails and/or text messages from CVS Specialty® about your prescription(s), account, and health care. Standard data rates apply. Message frequency varies. If unable to contact via text or email, Specialty Pharmacy will attempt to contact by phone. Primary Phone: ______ Alternate Phone: ______ Email: ______ Last Four of SSN: _____ Primary Language: ______ Parent/Caregiver/Legal Guardian Name (Last, First): _______Relationship to patient: ______ 2 PRESCRIBER INFORMATION Prescriber's Name: ______ DEA #: _____ Group or Hospital: ____ _____ State License #: ______ Address: _____ City, State, ZIP Code: _____ Contact's Phone: _____ Contact Person: ____ Contact's Phone: _____ INSURANCE INFORMATION Please fax copy of prescription and insurance cards with this form, if available (front and back) Is the Patient Insured? ☐ Yes ☐ No Is the Patient enrolled or eligible for Medicare/Medicaid? ☐ Yes ☐ No Policy Holder's Name:______ Policy Holder's DOB:______ Relationship to Patient:_____ Medical Insurance:
______ Policy ID:
_____ Group #:

Prescription Insurance:
______ Prescription Plan Telephone: _____ Group #: _____ RX BIN #: _____ RX PCN #:____ Policy ID: ☐ Check box if patient is enrolled in manufacturer copay assistance If yes, please provide ID# 4 DIAGNOSIS AND CLINICAL INFORMATION Needs by Date: Ship to: Patient Office Other: **Diagnosis (ICD-10):** Date of Diagnosis __/__/__ K50.00 Crohn's Disease of Small Intestine Without Complications K51.90 Ulcerative colitis, unspecified, without complications L40.50 Arthropathic Psoriasis, Unspecified L40.54 Juvenile Psoriatic Arthritis (JPsA) M06.9 Rheumatoid Arthritis, Unspecified M08.00 Juvenile Idiopathic Arthritis (JIA) M08.90 Polyarticular Juvenile Idiopathic Arthritis (PJIA) M08.20 Systemic Juvenile Idiopathic Arthritis (SJIA) M31.6 Giant Cell Arteritis (GCA) M32.1 Systemic lupus erythematosus (SLE) M32.14 Glomerular disease in systemic lupus erythematosus M45.9 Ankylosing Spondylitis of Unspecified Sites in Spine M45.A0 Non-Radiographic Axial Spondylarthritis (nr-axSpA) Other Code: ______Description: _____ Patient Clinical Information: □ NKDA Weight: □ kg □ lb Height: □ cm □ in Allergies: Treatment status: New to therapy Continuation of therapy; Date of last treatment __/__/__ TB Test Date __/__/ Positive Negative Hepatitis status: _____ Prior therapy, treatment dates, and reason(s) for discontinuation: **Nursing and Administration:** First dose administration of monoclonal antibodies (mABs) should be administered in a controlled setting (may vary depending upon medication specific policy). For Remicade/Remicade Biosimilars, the first dose must be administered in a controlled setting. Specialty pharmacy to coordinate home health Infusion nurse visit as necessary? Tyes No Site of Care: Home Infusion* Coram Ambulatory Infusion Suite (AIS)* Prescriber's Office** Other Infusion Clinic *Home Infusion/Coram AIS: Diluents, Flushes, Supplies, Nursing Services for drug administration/therapy teach train.

**Prescriber's Office/Other Infusion Clinic: Drug only for facility administration

Phone: 1-808-254-2727

NCPDP: 1203417

		Please Complete Patient and	Prescriber Information	
Patient Name:			Patient Phone:	
Prescriber Nam	ne:	P	rescriber Phone:	
Patient Clinica		_		
Allergies:		\square NKDA W. \square Continuation of therapy; \square	/eight: 🗌 kg 🗌 lb Height: 🗌 c	m 🗌 in
	// Positive		is status:	
Prior therapy, tre	eatment dates, and re	eason(s) for discontinuation:		
DDESCRIPT	ION INFORMATIO) N		
MEDICATION	STRENGTH		& DIRECTIONS	QUANTITY/REFILLS
☐ Actemra	80 mg/4 mL 200 mg/10 mL 400 mg/20 mL	☐ Induction Dose: Infuse 4 mg/kg eve	•	Quantity: Refills:
☐ Avsola	100 mg vial	(Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter		Quantity: # of 100 mg vial(s) Refills:
Benlysta	☐ 120 mg 5 mL vial ☐ 400 mg 20 mL vial	Induction Dose: 10 mg/kg IV (Dose =mg) at 2-week intervals for the first 3 doses and at 4-week intervals thereafter. Infuse IV over 1 hour.		Quantity: vials Refills:
☐ Entyvio	300 mg in a single dose vial in individual carton	☐ Induction Dose: 300 mg infused IV over 30 minutes at 0, 2 and 6 weeks, then every 8 weeks thereafter ☐ Maintenance Dose: 300 mg infused IV over 30 minutes every 8 weeks		Quantity: Refills:
Other	Strength:	□ Dose:		Quantity: Refills:
6 PRESCRIE	ER SIGNATURE	REQUIRED (STAMP SIGNAT	URE NOT ALLOWED)	
"Dispense As Written" / Brand Medically Necessary / Do Not Substitute / No Substitution / DAW / May Not Substitute		_	May Substitute / Product Selection Permitted / Substitution Permissible	D
Prescriber's S	Signature:	Date:	Prescriber's Signature:	Date:
CA, MA, NC & PR:	Interchange is mandated unle	ess Prescriber writes the words "No Substitution"	ATTN: New York and Iowa providers, pleas	e submit electronic prescription

Patient Name: _			
		Patient DOB: Patient Phone:	
Patient Address	:		
		Prescriber Phone:	
Patient Clinical			
Allergies:	n: Now to thoron	NKDA Weight: kg lb Height: lo Gontinuation of therapy; Date of last treatment//	cm ∐ in
	//_		
		eason(s) for discontinuation:	
	N INFORMATION	cason(s) for also on an action.	
MEDICATION		DOSE & DIRECTIONS	QUANTITY/REFILLS
		☐ Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg	(0.1
☐ Inflectra	100 mg vial	(Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter	Quantity: # of 100 mg vial(s) Refills:
Omvoh	300 mg/15 mL single dose vial	mg/kg (Dose =mg) every 8 weeks Induction Dose Week 0: Infuse 300 mg via IV infusion over at least 30 minutes Week 4: Infuse 300 mg via IV infusion over at least 30 minutes Week 8: Infuse 300 mg via IV infusion over at least 30 minutes	Quantity: Refills: 0 1 Vial 2 Vials 3 Vials
Orencia	250 mg vial	☐ Infuse mg at weeks 0, 2 and 4, then every 4 weeks thereafter	Quantity: Refills:
☐ Remicade	100 mg vial	Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one) Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: # of 100 mg vial(s) Refills:
Other	Strength:	Dose:	Quantity: Refills:
PRESCRIBER	SIGNATURE REOU	IIRED (STAMP SIGNATURE NOT ALLOWED)	
	-		
DAW / May Not Sub	•	essary / Do Not Substitute / No Substitution / May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:

	Pleas	se Complete Patient and I	Prescriber Information	
Patient Name: _		Patient DOB:	Patient Phone:	
Patient Address	:			
Prescriber Name		Pr	rescriber Phone:	
Patient Clinical				
Allergies:		UNKDA W	/eight: 🗌 kg 🗌 lb Height:	
Treatment status	s: New to therapy	☐ Continuation of therapy; D	eate of last treatment//	
			s status:	
	ION INFORMATION) for discontinuation:		
MEDICATION	STRENGTH	DOSE	& DIRECTIONS	QUANTITY/REFILLS
Riabni			a DIRECTIONS	QUANTITY REFIELD
Rituxan Ruxience	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000	mg separated by 2 weeks	Quantity: Refills:
Saphnelo	300 mg/2 mL (150 mg/mL)	300 mg IV over a 30-mir	nute period, every 4 weeks	Quantity: vials Refills:
		☐ Week 4: Infuse 2 mg/kg	IV (Dose=mg) over 30 minutes IV (Dose=mg) over 30 minutes	Quantity: vials Refills: 0 Quantity: vials Refills: 0
Simponi	50 mg/4 mL single dose vial	weeks	=mg) over 30 minutes every 8	Quantity: vials Refills:
ARIA	dose viai	minutes	old) Induction Dose n² IV (Dose=mg) over 30 n² IV (Dose=mg) over 30	Quantity: vials Refills: 0 Quantity: vials Refills: 0
		8 weeks	old) Maintenance Dose se=mg) over 30 minutes every	Quantity: vials Refills:
Skyrizi	600 mg/10 mL (60 mg/mL) single dose vial	Week 4: Infuse 1,200 mg	V over at least one hour	Quantity: 1 vial Refills: 0 Quantity: 1 vial Refills: 0 Quantity: 1 vial Refills: 0 Quantity: 2 vials Refills: 0 Quantity: 2 vials Refills: 0 Quantity: 2 vials Refills: 0 Refills: 0 Refills: 0
☐ Stelara	130 mg/26 mL (5 mg/mL) IV single- dose vial	more than 55 kg to 85 kg used 3	week 0: # of vials to be used 2 g 390 mg at week 0: # of vials to be g at week 0: # of vials to be used 4	Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
6 PRESCRIB	ER SIGNATURE REQU	JIRED (STAMP SIGNAT	URE NOT ALLOWED)	
	en" / Brand Medically Necessary / Do pstitute	_	May Substitute / Product Selection Permitted / Substitution Permissible Prescriber's Signature:	Date:
CA. MA. NC & PR:	nterchange is mandated unless Prescri	iber writes the words "No Substitution"	ATTN: New York and Iowa provide	rs, please submit electronic prescription

	Pleas	se Complete Patient and Prescriber Information	
Patient Name: _		Patient DOB: Patient Phone:	
Patient Address	:		
Prescriber Nam	e:	Prescriber Phone:	
Patient Clinical	Information:		
Allergies:		NKDA Weight: 🗌 kg 🗌 lb Height:	cm 🗌 in
Treatment status	s: New to therapy	Continuation of therapy; Date of last treatment//	
		egative Hepatitis status:	
) for discontinuation:	<u> </u>
	ION INFORMATION		
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS
☐ Tremfya	200 mg/20 mL (10 mg/mL) single- dose vial	Intravenous UC or CD Induction Dose: Week 0: Infuse 200 mg IV over at least one hour Week 4: Infuse 200 mg IV over at least one hour Week 8: Infuse 200 mg IV over at least one hour	Quantity: 1 Vial Refills: 0 Quantity: 1 Vial Refills: 0 Quantity: 1 Vial Refills: 0
☐ Truxima	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separated by 2 weeks ☐ Other:	Quantity: Refills:
Tyenne (tocilizumab- aazg)	☐ 80 mg/4 mL vial ☐ 200 mg/10 mL vial ☐ 400 mg/20 mL vial	RA Induction Dose: Infuse 4 mg per kg (mg) IV every 4 weeks RA Maintenance Dose: Infuse 8 mg per kg (mg) IV every 4 weeks (doses exceeding 800 mg per infusion are not recommended) Giant Cell Arteritis Dose: Infuse 6 mg per kg (mg) IV every 4 weeks (doses exceeding 600 mg per infusion are not recommended) PJIA Dose (≥ 2 years old weighing < 30 kg): Infuse 10 mg per kg (mg) IV every 4 weeks PJIA Dose (≥ 2 years old weighing ≥ 30 kg): Infuse 8 mg per kg (mg) IV every 4 weeks SJIA Dose (≥ 2 years old weighing < 30 kg): Infuse 12 mg per kg (mg) IV every 2 weeks SJIA Dose (≥ 2 years old weighing ≥ 30 kg): Infuse 8 mg per kg (mg) IV every 2 weeks Other:	Quantity: (# of 80 mg vials) (# of 200 mg vials) (# of 400 mg vials) Refills:
Other	Strength:	☐ Dose:	Quantity: Refills:
"Dispense As Writt DAW / May Not Sul Prescriber's S	en" / Brand Medically Necessary / D ostitute ignature:	Substitution Permissible Prescriber's Signature:	

Autoimmune IV Enrollment Form Nursing Orders

	Pleas	se Complete Patient and Prescriber Inforn	nation
		Patient DOB: P	
Patient Address:			
Prescriber Name:		Prescriber Phone:	
Patient Clinical Informati			
Allergies:		NKDA Weight: kg	
reatment status: New t		Continuation of therapy; Date of last treatment	
B Test Date//[
PRESCRIPTION INFO) for discontinuation:	DE INTERCORDE DONE AT HOME (OOD AM AIGH
MEDICATION/SUPPLIES	ROUTE	**ITEMS BELOW THIS LINE WILL ONLY BE SENT FO DOSE /STRENGTH/ DIRECTIONS	
WEDICATION/SUPPLIES	ROUTE	Catheter Care/Flush - Only on drug admin days -	
		maintain IV access and patency	SAGIT OF TRIVES
Catheter:		PIV: NS 5 mL (Heparin 10 units/mL 3-5 mL if multip	ple days) Quantity:
☐ PIV ☐ PORT	IV	CVC/PICC: NS 10 mL & Heparin 10 units/mL or	
☐ CVC/PICC		3-5 mL.	
		PORT: 10 mL sterile saline to access PORT w/ hub	er needle
		NS 10 mL & Heparin 100 units/mL 3-5mL.	
		- C C C	Hydration max infusion
Hydration:	15.7	Pre: ☐ 500 mL ☐ 1000 mL ☐ 0ther:	rate mL/hr
☐ NS ☐ D5W	IV	Concurrent: ☐ 500 mL ☐ 1000 mL ☐ Other: Post: ☐ 500 mL ☐ 1000 mL ☐ Other:	
		Post: [] 500 mL [] 1000 mL [] Other:	otherwise indicated)
		1:1000, 0.3mg/0.3 mL (greater than 30 kg/66 l	,
		1:1000, 0:311g/0:311L (greater trial 30 kg/30 to 1:1000, 0:15mg/0:311L (15-30 kg/33-66 lbs)	
Epinephrine	□ ім	1:1000, 0.1 mg/kg, Max 0.3mg (under 15kg)	Quantity:
nursing requires	□ sc	Mild-Moderate Reactions. May repeat in 3-5 minu	utes as needed Refills:
		for severe allergic reaction, also call 911	
Diphenhydramine		Premedication:	
Oral	PO	☐ 12.5 mg/kg (0-30 kg)	Quantity:
Orai		25 mg	Refills:
		50 mg (Over 30 kg)	
		1 mg/kg (under 15 kg)	
Diphenhydramine		12.5 mg-50 mg (15-30 kg)	
50 mg/mL vial	Slow IV	25 mg-50 mg (Over 30 kg)	Quantity:
nursing required	□ ім	If mild/moderate reaction: may repeat in 3-5 minu (Adult max dose: 100 mg/day)	ites as needed Refills:
		If severe allergic reaction: call 911	
	Peripheral		
	Access	10 mL NS post flush	Send quantity
Flush Orders:	Central	50 mL NS post flush to clear medication from t	<u> </u>
	Venous	(recommended if no post-hydration)	for medication days
	Access	Other:	supply
Additional	-		
Medication:			
Patient is interested in patient supp PRESCRIBER SIGN		STAMP SIGNATURE NOT ALLOWED JIRED (STAMP SIGNATURE NOT ALLO)	Ancillary supplies and kits provided as needed for administratio
		•	-
"Dispense As Written" / Brand Me DAW / May Not Substitute	calcally Necessary / De	o Not Substitute / No Substitution / May Substitute / Product Substitution Permissible	
		Date: Prescriber's Signa	ature:Date:

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Specialty and/or one of its affiliates.