

Fax Referral To: 1-855-297-1270 Phone: 1-888-280-1190

Address: 6020 Ave Roberto Sanchez Vilella Carolina, PR 00982 NCPDP: 4026325

Six Simple S	Steps to Subm	itting a Referra	ıl	
PATIENT INFORMATION (Complete or include de	lemographic she			
Patient Name:		DOB:		Gender: Male Female
Address:		•		
Preferred Contact Methods:  Phone (to primary # prov below)	vided below) 🔲	Text (to cell # pro	ovided below)	Email (to email provided
Note: Carrier charges may apply. By providing the phone	number(s) and e	email address abo	ve, you are co	nsenting to receive
automated calls, emails and/or text messages from CVS S				
rates apply. Message frequency varies. If unable to contac	ct via text or em	ail, Specialty Phar	macy will atte	mpt to contact by phone.
Primary Phone:	A	Alternate Phone: _		
Email:				
Parent/Caregiver/Legal Guardian Name (Last, First):		_Relationship to	patient:	
2 PRESCRIBER INFORMATION				
Prescriber's Name:		State License #		
Prescriber's Name: Group or Hos	 nital:	_Otato Elocitoo III.		
Address:	City. S	tate. ZIP Code:		
Address: Fax C	contact Person:		Contact's Ph	one:
3 INSURANCE INFORMATION Please fax copy of plates the Detion to present on the Detion to				
Is the Patient Insured? Yes No Is the Patient enro				
Policy Holder's Name: Tele Medical Insurance: Tele				
Prescription Insurance: Group #:		_ Prescription Pta	ın retephone: . -	
☐ Check box if patient is enrolled in manufacturer copay				
_	-	es, piease providi	- ID#	
4 DIAGNOSIS AND CLINICAL INFORMATION				
Needs by Date: Ship to: Datient Ship to: Ship to: Ship to: Ship to: Datient Ship to: Sh	Office 🗌 Other	r:		
Diagnosis (ICD-10): Date of Diagnosis//	/			
K50.00 Crohn's Disease of Small Intestine Without Co				
K51.90 Ulcerative colitis, unspecified, without complic	•			
L40.50 Arthropathic Psoriasis, Unspecified				
L40.54 Juvenile Psoriatic Arthritis (JPsA)				
M06.9 Rheumatoid Arthritis, Unspecified				
M08.00 Juvenile Idiopathic Arthritis (JIA)				
M08.90 Polyarticular Juvenile Idiopathic Arthritis (PJI.	A)			
M08.20 Systemic Juvenile Idiopathic Arthritis (SJIA)				
M31.6 Giant Cell Arteritis (GCA)				
M32.1 Systemic lupus erythematosus (SLE)				
M32.14 Glomerular disease in systemic lupus erythem				
M45.9 Ankylosing Spondylitis of Unspecified Sites in S	Spine			
M45.A0 Non-Radiographic Axial Spondylarthritis (nr-a	axSpA)			
Other Code:Description:	:			
Patient Clinical Information:				
_	IKDA We	ight: 🗌 kg	☐ lb Height: _	🗌 cm 🔲 in
Allergies: New to therapy Continuation	on of therapy; Da	te of last treatmen	t//	
TB Test Date// Positive Negative Prior therapy, treatment dates, and reason(s) for discontinua	☐ Hepatitis	status:		
	ation:			
Nursing and Administration:				
First dose administration of monoclonal antibodies (mAB	s) should be adr	ministered in a co	ntrolled setting	g (may vary depending upon
medication specific policy).				
For Remicade/Remicade Biosimilars, the first dose mu				
Specialty pharmacy to coordinate home health Infusion r				
Site of Care: Home Infusion* Coram Ambulatory				
*Home Infusion/Coram AIS: Diluents, Flushes, Supplies,			stration/therap	by teach train.
**Prescriber's Office/Other Infusion Clinic: Drug only for	facility administ	ration		

		Please Complete Patient and	Prescriber Information	
Patient Name:			Patient Phone:	
Prescriber Nam	ne:	P	rescriber Phone:	
Patient Clinica		_		
Allergies:		$\square$ NKDA W. $\square$ Continuation of therapy; $\square$	/eight: 🗌 kg 🗌 lb Height: 🗌 c	m 🗌 in
	_//_ Positive		is status:	
Prior therapy, tre	eatment dates, and re	eason(s) for discontinuation:		
DDESCRIPT	ION INFORMATIO	) N		
MEDICATION	STRENGTH		& DIRECTIONS	QUANTITY/REFILLS
☐ Actemra	80 mg/4 mL 200 mg/10 mL 400 mg/20 mL	☐ Induction Dose: Infuse 4 mg/kg eve	•	Quantity: Refills:
☐ Avsola	100 mg vial	5 mg/kg (Dose =mg) at week  Crohn's Disease (Adult) Maintenand (Dose =mg) every 8 weeks  Crohn's Disease (Pediatric ≥ 6 year Infuse IV at 5 mg/kg (Dose =r  Plaque Psoriasis & Psoriatic Arthriti (Dose =mg) at weeks 0, 2, 6 at Plaque Psoriasis & Psoriatic Arthriti Infuse IV at 5 mg/kg (Dose =r  Rheumatoid Arthritis Induction Dos (Dose =mg) at weeks 0, 2, 6 and Rheumatoid Arthritis Maintenance (Dose =mg) every 4, 6 or 8 we Ulcerative Colitis (Adult and Pediatis 5 mg/kg (Dose =mg) at week	every 6 weeks thereafter  e Dose: Infuse IV at 5 mg/kg  ic ≥ 6 years old) Induction Dose: Infuse IV at s 0, 2, 6 and every 8 weeks thereafter  ce Dose: Infuse IV at 5-10 mg/kg  s old) Maintenance Dose: mg) every 8 weeks is Induction Dose: Infuse IV at 5 mg/kg and every 8 weeks thereafter is Maintenance Dose: mg) every 8 weeks thereafter is Maintenance Dose: mg) every 8 weeks  ie: Infuse IV at 3 mg/kg every 8 weeks thereafter Dose: Infuse IV at 3-10 mg/kg eks (circle one) ric ≥ 6 years old) Induction Dose: Infuse IV at s 0, 2, 6 and every 8 weeks thereafter ric ≥ 6 years old) Maintenance Dose: Infuse IV	Quantity: # of 100 mg vial(s) Refills:
Benlysta	☐ 120 mg 5 mL vial ☐ 400 mg 20 mL vial	Induction Dose: 10 mg/kg IV (Dose =mg) at 2-week intervals for the first 3 doses and at 4-week intervals thereafter. Infuse IV over 1 hour.		Quantity: vials Refills:
☐ Entyvio	300 mg in a single dose vial in individual carton	☐ Induction Dose: 300 mg infused IV over 30 minutes at 0, 2 and 6 weeks, then every 8 weeks thereafter ☐ Maintenance Dose: 300 mg infused IV over 30 minutes every 8 weeks		Quantity: Refills:
Other	Strength:	□ Dose:		Quantity: Refills:
6 PRESCRIE	ER SIGNATURE	REQUIRED (STAMP SIGNAT	URE NOT ALLOWED)	
DAW / May Not Su	bstitute	ssary / Do Not Substitute / No Substitution /	May Substitute / Product Selection Permitted / Substitution Permissible	<b>D</b>
Prescriber's S	Signature:	Date:	Prescriber's Signature:	Date:
CA, MA, NC & PR:	Interchange is mandated unle	ess Prescriber writes the words "No Substitution"	ATTN: New York and Iowa providers, pleas	e submit electronic prescription

Patient Name: _			
		Patient DOB: Patient Phone:	
Patient Address	<b>:</b>		
		Prescriber Phone:	
Patient Clinical			
Allergies:	n: Now to thoron	NKDA Weight: kg  lb Height: lo Gontinuation of therapy; Date of last treatment//	cm ∐ in
	//_		
		eason(s) for discontinuation:	
	N INFORMATION	cason(s) for also on an action.	
MEDICATION		DOSE & DIRECTIONS	QUANTITY/REFILLS
		☐ Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg	(0.1
☐ Inflectra	100 mg vial	(Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter  Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks  Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks  Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks  Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks  Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one)  Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter	Quantity: # of 100 mg vial(s) Refills:
Omvoh	300 mg/15 mL single dose vial	mg/kg (Dose =mg) every 8 weeks  Induction Dose  Week 0: Infuse 300 mg via IV infusion over at least 30 minutes  Week 4: Infuse 300 mg via IV infusion over at least 30 minutes  Week 8: Infuse 300 mg via IV infusion over at least 30 minutes	Quantity: Refills: 0  1 Vial 2 Vials 3 Vials
Orencia	250 mg vial	☐ Infuse mg at weeks 0, 2 and 4, then every 4 weeks thereafter	Quantity: Refills:
☐ Remicade	100 mg vial	Ankylosing Spondylitis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 6 weeks thereafter  Ankylosing Spondylitis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 6 weeks  Crohn's Disease (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Crohn's Disease (Adult) Maintenance Dose: Infuse IV at 5-10 mg/kg (Dose =mg) every 8 weeks  Crohn's Disease (Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks  Plaque Psoriasis & Psoriatic Arthritis Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Plaque Psoriasis & Psoriatic Arthritis Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks  Rheumatoid Arthritis Induction Dose: Infuse IV at 3 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Rheumatoid Arthritis Maintenance Dose: Infuse IV at 3-10 mg/kg (Dose =mg) every 4, 6 or 8 weeks (circle one)  Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Induction Dose: Infuse IV at 5 mg/kg (Dose =mg) at weeks 0, 2, 6 and every 8 weeks thereafter  Ulcerative Colitis (Adult and Pediatric ≥ 6 years old) Maintenance Dose: Infuse IV at 5 mg/kg (Dose =mg) every 8 weeks	Quantity: # of 100 mg vial(s) Refills:
Other	Strength:	Dose:	Quantity: Refills:
PRESCRIBER	SIGNATURE REOU	IIRED (STAMP SIGNATURE NOT ALLOWED)	
	-		
DAW / May Not Sub	•	essary / Do Not Substitute / No Substitution /  May Substitute / Product Selection Permitted / Substitution Permissible  Prescriber's Signature:	Date:

	Pleas	se Complete Patient and I	Prescriber Information	
Patient Name: _		Patient DOB:	Patient Phone:	
Patient Address	:			
Prescriber Name		Pr	rescriber Phone:	
Patient Clinical				
Allergies:		UNKDA W	/eight: 🗌 kg 🗌 lb Height:	
Treatment status	s: New to therapy	☐ Continuation of therapy; D	eate of last treatment//	
			s status:	<del></del>
	ION INFORMATION	) for discontinuation:		
MEDICATION	STRENGTH	DOSE	& DIRECTIONS	QUANTITY/REFILLS
Riabni			a DIRECTIONS	QUANTITY REFIELD
Rituxan Ruxience	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000	mg separated by 2 weeks	Quantity: Refills:
Saphnelo	300 mg/2 mL (150 mg/mL)	300 mg IV over a 30-mir	nute period, every 4 weeks	Quantity: vials Refills:
		☐ Week 4: Infuse 2 mg/kg	IV (Dose=mg) over 30 minutes IV (Dose=mg) over 30 minutes	Quantity: vials Refills: 0 Quantity: vials Refills: 0
Simponi	50 mg/4 mL single dose vial	weeks	=mg) over 30 minutes every 8	Quantity: vials Refills:
ARIA dose viai	minutes	old) Induction Dose  n² IV (Dose=mg) over 30  n² IV (Dose=mg) over 30	Quantity: vials Refills: 0 Quantity: vials Refills: 0	
		8 weeks	old) Maintenance Dose se=mg) over 30 minutes every	Quantity: vials Refills:
Skyrizi	600 mg/10 mL (60 mg/mL) single dose vial	Week 4: Infuse 1,200 mg	V over at least one hour	Quantity: 1 vial Refills: 0 Quantity: 1 vial Refills: 0 Quantity: 1 vial Refills: 0 Quantity: 2 vials Refills: 0 Quantity: 2 vials Refills: 0 Quantity: 2 vials Refills: 0 Refills: 0 Refills: 0
☐ Stelara	130 mg/26 mL (5 mg/mL) IV single- dose vial	more than 55 kg to 85 kg used 3	week 0: # of vials to be used 2 g 390 mg at week 0: # of vials to be g at week 0: # of vials to be used 4	Quantity: 2 Vials 3 Vials 4 Vials Refills: 0
6 PRESCRIB	ER SIGNATURE REQU	JIRED (STAMP SIGNAT	URE NOT ALLOWED)	
	en" / Brand Medically Necessary / Do pstitute	_	May Substitute / Product Selection Permitted / Substitution Permissible  Prescriber's Signature:	Date:
CA. MA. NC & PR:	nterchange is mandated unless Prescri	iber writes the words "No Substitution"	ATTN: New York and Iowa provide	rs, please submit electronic prescription

	Pleas	se Complete Patient and Prescriber Information	
Patient Name: _		Patient DOB: Patient Phone:	
Patient Address	<b>:</b>		
Prescriber Nam	e:	Prescriber Phone:	
Patient Clinical	Information:		
Allergies:		NKDA Weight: 🗌 kg 🗌 lb Height:	cm 🗌 in
Treatment status	s: New to therapy	Continuation of therapy; Date of last treatment//	
		egative  Hepatitis status:	
		) for discontinuation:	<u> </u>
	ION INFORMATION		
MEDICATION	STRENGTH	DOSE & DIRECTIONS	QUANTITY/REFILLS
☐ Tremfya	200 mg/20 mL (10 mg/mL) single- dose vial	Intravenous UC or CD Induction Dose:  Week 0: Infuse 200 mg IV over at least one hour Week 4: Infuse 200 mg IV over at least one hour Week 8: Infuse 200 mg IV over at least one hour	Quantity: 1 Vial Refills: 0 Quantity: 1 Vial Refills: 0 Quantity: 1 Vial Refills: 0
☐ Truxima	☐ 100 mg/10 mL vial ☐ 500 mg/50 mL vial	☐ Infuse two doses of 1000 mg separated by 2 weeks ☐ Other:	Quantity: Refills:
Tyenne (tocilizumab- aazg)	☐ 80 mg/4 mL vial ☐ 200 mg/10 mL vial ☐ 400 mg/20 mL vial	RA Induction Dose: Infuse 4 mg per kg ( mg) IV every 4 weeks  RA Maintenance Dose: Infuse 8 mg per kg ( mg) IV every 4 weeks (doses exceeding 800 mg per infusion are not recommended)  Giant Cell Arteritis Dose: Infuse 6 mg per kg ( mg) IV every 4 weeks (doses exceeding 600 mg per infusion are not recommended)  PJIA Dose ( ≥ 2 years old weighing < 30 kg): Infuse 10 mg per kg ( mg) IV every 4 weeks  PJIA Dose ( ≥ 2 years old weighing ≥ 30 kg): Infuse 8 mg per kg ( mg) IV every 4 weeks  SJIA Dose ( ≥ 2 years old weighing < 30 kg): Infuse 12 mg per kg ( mg) IV every 2 weeks  SJIA Dose ( ≥ 2 years old weighing ≥ 30 kg): Infuse 8 mg per kg ( mg) IV every 2 weeks  Other:	Quantity:  (# of 80 mg vials)  (# of 200 mg vials)  (# of 400 mg vials)  Refills:
Other	Strength:	☐ Dose:	Quantity: Refills:
"Dispense As Writt DAW / May Not Sul Prescriber's S	en" / Brand Medically Necessary / D ostitute ignature:	Substitution Permissible Prescriber's Signature:	

## Autoimmune IV Enrollment Form Nursing Orders

	Pleas	se Complete Patient and Prescriber Informat	tion
		Patient DOB: Pati	
Patient Address:			
Prescriber Name:		Prescriber Phone:	
Patient Clinical Informati	<del></del>		
Allergies:		NKDA Weight: kg	
reatment status: New		Continuation of therapy; Date of last treatment	
B Test Date//			
PRESCRIPTION INFO		for discontinuation:	UNITED CONTRACT LIGHT (COD AND ALCOU
MEDICATION/SUPPLIES	ROUTE	**ITEMS BELOW THIS LINE WILL ONLY BE SENT FOR DOSE /STRENGTH/ DIRECTIONS	QUANTITY/REFILLS
WEDICATION/SUPPLIES	ROOTE	Catheter Care/Flush - Only on drug admin days - SA	-
		maintain IV access and patency	SIT OF FIXIVES
Catheter:		PIV: NS 5 mL (Heparin 10 units/mL 3-5 mL if multiple	days) Quantity:
☐ PIV ☐ PORT	IV	CVC/PICC: NS 10 mL & Heparin 10 units/mL or	
☐ CVC/PICC		3-5 mL.	
		PORT: 10 mL sterile saline to access PORT w/ huber	needle
		NS 10 mL & Heparin 100 units/mL 3-5mL.	
		- U : U : U	Hydration max infusion
Hydration:		Pre: ☐ 500 mL ☐ 1000 mL ☐ Other:	rate mL/hr
☐ NS ☐ D5W	IV	Concurrent: ☐ 500 mL ☐ 1000 mL ☐ Other: Post: ☐ 500 mL ☐ 1000 mL ☐ Other:	
		Post:	otherwise indicated)
		1:1000, 0.3mg/0.3 mL (greater than 30 kg/66 lbs)	,
		1:1000, 0.511g/0.5 HL (greater trial 30 kg/30 lbs)	
☐ <i>Epinephrine</i>	□ ім	1:1000, 0.1 mg/kg, Max 0.3mg (under 15kg)	Quantity:
**nursing requires**	□sc	Mild-Moderate Reactions. May repeat in 3-5 minutes	s as needed Refills:
		for severe allergic reaction, also call 911	
Diphenhydramine		Premedication:	
Oral	PO	12.5 mg/kg (0-30 kg)	Quantity:
Orai		25 mg	Refills:
		50 mg (Over 30 kg)	
		1 mg/kg (under 15 kg)	
Diphenhydramine		12.5 mg-50 mg (15-30 kg)	
50 mg/mL vial	Slow IV	25 mg-50 mg (Over 30 kg)	Quantity:
**nursing required**	□ ім	If mild/moderate reaction: may repeat in 3-5 minutes (Adult max dose: 100 mg/day)	s as needed Refills:
		If severe allergic reaction: call 911	
	Peripheral		
	Access	10 mL NS post flush	Send quantity
Flush Orders:	Central	50 mL NS post flush to clear medication from tub	_
	Venous	(recommended if no post-hydration)	for medication days
	Access	Other:	supply
Additional			
Medication:			
Patient is interested in patient supp	. •	STAMP SIGNATURE NOT ALLOWED  JIRED (STAMP SIGNATURE NOT ALLOW	illary supplies and kits provided as needed for administration
	· ·		•
•	eaically Necessary / D	o Not Substitute / No Substitution / May Substitute / Product Se Substitution Permissible	election Permitted /
DAW / May Not Substitute			
DAW / May Not Substitute Prescriber's Signature: _			re:Date:

The information provided above is true and accurate to the best of my knowledge, with supporting documentation in the patient's medical record. By signing above, I hereby authorize CVS Specialty Pharmacy and/or its affiliate pharmacies to complete and submit prior authorization (PA) requests to payors for the prescribed medication for this patient and to attach this Enrollment Form to the PA request as my signature.

CONFIDENTIALITY NOTICE: This communication and any attachments may contain confidential and/or privileged information for the use of the designated recipients named above. If you are not the intended recipient, you are hereby notified that you have received this communication in error and that any review, disclosure, dissemination, distribution or copying of it or its contents is prohibited. If you have received this communication in error, please notify the sender immediately by telephone and destroy all copies of this communication and any attachments.

Plan member privacy is important to us. Our employees are trained regarding the appropriate way to handle members' private health information.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Specialty and/or one of its affiliates.