



Visit cvs.co/ai-list or scan the QR code to view electronically.
For easy access, you can bookmark this page.

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Actemra (tocilizumab)	Genentech	<ul style="list-style-type: none"> • CRS (2+) • GCA • PJIA (2+) • RA • SJIA (2+) • SSc-ILD • Hospitalized adults who have COVID-19 	SC injection: <ul style="list-style-type: none"> • Self-administered IV infusion: <ul style="list-style-type: none"> • Prepared and infused by a health care professional 	SC: <ul style="list-style-type: none"> • 162 mg/0.9 mL PFS • 162 mg/0.9 mL PEN IV: <ul style="list-style-type: none"> • 80 mg/4 mL • 200 mg/10 mL • 400 mg/20 mL 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light • Keep syringes dry 	Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions
Adalimumab-aacf (unbranded version of Idacio)	Fresenius Kabi	<ul style="list-style-type: none"> • AS • CD (6+) • HS (12+) • PJIA (2+) • PsA • PsO • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Adalimumab-aaty (unbranded version of Yuflyma)	Celltrion USA	SC injection: <ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsA • PsO • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 40 mg/0.4 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Adalimumab-adaz (unbranded version of Hyrimoz)	Sandoz	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 10 mg/0.1 mL PFS • 20 mg/0.2 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.4 mL PEN • 80 mg/0.8 mL PEN 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Adalimumab-fkjp (unbranded version of Huloio)	Mylan	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS • 20 mg/0.4 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original package 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Adbry (tralokinumab-ldrm)	LEO Pharma	<ul style="list-style-type: none"> • AD (12+) 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 150 mg/mL PFS • 300 mg/2 mL PEN 	<ul style="list-style-type: none"> • Store refrigerated at 36°F to 46°F (2°C to 8°C) • Store in the original carton to protect from light 	Upper respiratory tract infections, conjunctivitis, injection site reactions and eosinophilia

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Amjevita (adalimumab-atto)	Amgen	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 80 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.4 mL PFS • 20 mg/0.2 mL PFS • 10 mg/0.2 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original package 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Arcalyst (rilonacept)	Kiniksa Pharmaceuticals	<ul style="list-style-type: none"> • CAPS (12+) • FCAS (12+) • MWS (12+) • DIRA (≥10 kg) • RP (12+) 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 220 mg of lyophilized powder in a single dose vial for reconstitution 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in original carton • Protect from light • Do not use beyond the date stamped on the label • After reconstitution, may be kept at room temperature, protected from light 	Injection-site reactions and upper respiratory tract infections

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Avsola (infliximab-axxq)	Amgen	<ul style="list-style-type: none"> AS CD (6+) PsO PsA RA UC (6+) 	IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg lyophilized powder in a single dose vial for reconstitution 	<ul style="list-style-type: none"> Store unopened Avsola vials in a refrigerator at 36°F to 46°F (2°C to 8°C) Protect from light 	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Benlysta (belimumab)	GlaxoSmithKline	<ul style="list-style-type: none"> SLE (5+) LN (5+) 	IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional SC injection: <ul style="list-style-type: none"> Self-administered 	IV: <ul style="list-style-type: none"> 120 mg lyophilized powder in a single dose vial for reconstitution 400 mg lyophilized powder in a single dose vial for reconstitution SC: <ul style="list-style-type: none"> 200 mg/mL PEN 200 mg/mL PFS 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Store in original packages Protect from light Avoid exposure to heat 	Nausea, diarrhea, pyrexia, nasopharyngitis, bronchitis, pharyngitis, insomnia, pain in extremity, depression and migraine
Bimzelx (bimekizumab-bkzx)	UCB	<ul style="list-style-type: none"> AS HS Nr-axSPA PsO PsA 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 160mg/mL PEN 160mg/mL PFS 320 mg/2 mL PEN 320 mg/2 mL PFS 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Do not shake Store in original carton to protect from light 	Upper respiratory tract infections, oral candidiasis, headache, injection site reactions, tinea infections, gastroenteritis, Herpes simplex infections, acne, folliculitis, other candida infections and fatigue

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Cibinqo (abrocitinib)	Pfizer	<ul style="list-style-type: none"> AD (12+) 	Oral	Oral: <ul style="list-style-type: none"> 50 mg tablets 100 mg tablets 200 mg tablets 	<ul style="list-style-type: none"> Store at 68°F to 77°F (20°C to 25°C) Keep in original package 	Nasopharyngitis, nausea, headache, herpes simplex, increased blood creatinine phosphokinase, dizziness, urinary tract infection, fatigue, acne, vomiting, oropharyngeal pain, influenza, gastroenteritis, impetigo, hypertension, contact dermatitis, upper abdominal pain, abdominal discomfort, herpes zoster and thrombocytopenia
Cimzia (certolizumab pegol)	UCB	<ul style="list-style-type: none"> AS CD Nr-axSpA PJIA (2+) PsO PsA RA 	SC injection: <ul style="list-style-type: none"> Self-administered (PFS only) 	SC: <ul style="list-style-type: none"> 200 mg/mL PFS 200 mg lyophilized powder in a single- dose vial (for health care professional use only) 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Protect from light 	Upper respiratory tract infection, rash and urinary tract infection
Cosentyx (secukinumab)	Novartis	SC: <ul style="list-style-type: none"> AS ERA (4+) HS Nr-axSpA PsO (6+) PsA (2+) 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 75 mg/0.5 mL PFS 150 mg/mL PEN 150 mg/mL PFS 300 mg/2 mL PEN 300 mg/2 mL PFS 	SC: <ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Keep the product in the original carton Protect from light Do not shake to avoid foaming 	Nasopharyngitis, diarrhea and upper respiratory tract infection

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Dupixent (dupilumab)	Sanofi	<ul style="list-style-type: none"> • AD (6 mo+) • Asthma (6+) • COPD • CRSwNP (12+) • CRSwNP • EoE (1+ and ≥ 15 kg) • PN 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 100 mg/0.67 mL PFS • 200 mg/1.14 mL PEN • 200 mg/1.14 mL PFS • 300 mg/2 mL PEN • 300 mg/2 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light • Do not freeze 	Injection site reactions, conjunctivitis, blepharitis, oral herpes, keratitis, eye pruritus, other herpes simplex virus infection and dry eye
Ebglyss (lebrikizumab-lbkz)	Eli Lilly	<ul style="list-style-type: none"> • AD (12+) 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 250 mg/2 mL PFS • 250 mg/2 mL PEN 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in the original carton to protect from light • Do not freeze • Do not shake 	Conjunctivitis, injection site reactions, and herpes zoster

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Enbrel (etanercept)	Amgen	<ul style="list-style-type: none"> AS JPsA (2+) PsO (4+) PsA (2+) RA 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 25 mg/0.5 mL single dose vial 25 mg lyophilized powder in a multiple dose vial for reconstitution 25 mg/0.5 mL PFS 50 mg/mL PFS 50 mg/mL PEN 50 mg/mL Enbrel Mini single-dose prefilled cartridge 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Do not shake Store in the original carton to protect from light or physical damage 	Infections and injection site reactions
Entyvio (vedolizumab)	Takeda	<ul style="list-style-type: none"> CD UC 	IV infusion: <ul style="list-style-type: none"> Administered by a healthcare provider SC injection: <ul style="list-style-type: none"> Self-administered 	IV: <ul style="list-style-type: none"> 300 mg single dose vial SC: <ul style="list-style-type: none"> 108 mg/0.68 mL PEN 	<ul style="list-style-type: none"> Refrigerate unopened vials, PEN, and PFS at 36°F to 46°F (2°C to 8°C) Do not freeze Retain in original package to protect from light 	Nasopharyngitis, headache, arthralgia, nausea, pyrexia, upper respiratory tract infection, fatigue, cough, bronchitis, influenza, back pain, rash, pruritus, sinusitis, oropharyngeal pain and pain in extremities. In addition, injection site reactions are associated with the SC injection only.

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Hadlima (adalimumab-bwwd)	Samsung Bioepis/ Organon	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 40 mg/0.8 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PEN 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original container 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Hulio (adalimumab-fkjp)	Mylan	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS • 20 mg/0.4 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Humira (adalimumab)	AbbVie/ Cordavis	<ul style="list-style-type: none"> • AS • CD (6+) • HS (12+) • PJIA (2+) • PsO • PsA • RA • UC (5+) • UV (2+) 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 80 mg/0.8 mL PEN • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.8 mL PFS • 40 mg/0.4 mL PFS • 20 mg/0.2 mL PFS • 10 mg/0.1 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Hyrimoz (adalimumab-adaz)	Cordavis/ Sandoz	<ul style="list-style-type: none"> • AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 80 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PFS • 40 mg/0.8 mL PFS • 20 mg/0.2 mL PFS 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Idacio (adalimumab-aacf)	Fresenius Kabi	<ul style="list-style-type: none"> AS CD (6+) HS PJIA (2+) PsO PsA RA UC UV 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 40 mg/0.8 mL PEN 40 mg/0.8 mL PFS 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Protect from light 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Ilaris (canakinumab)	Novartis	<ul style="list-style-type: none"> CAPS (4+) FCAS (4+) FMF Gout flares HIDS/MKD MWS (4+) TRAPS Active Still's Disease/AOSD (2+) SJIA (2+) 	SC injection: <ul style="list-style-type: none"> Should only be administered by a health care professional 	SC: <ul style="list-style-type: none"> 150 mg/mL single dose vial 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Store in the original carton to protect from light 	<p>CAPS: Nasopharyngitis, diarrhea, influenza, rhinitis, nausea, headache, bronchitis, gastroenteritis, pharyngitis, weight increased, musculoskeletal pain and vertigo</p> <p>TRAPS, HIDS/MKD, and FMF: injection-site reactions and nasopharyngitis</p> <p>Still's Disease: Nasopharyngitis and upper respiratory tract infections, abdominal pain and injection-site reactions</p> <p>Gout Flares: Nasopharyngitis, upper respiratory tract infections, urinary tract infections, hypertriglyceridemia and back pain</p>

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Ilumya (tildrakizumab-asmn)	Sun Pharma	PsO	SC injection: <ul style="list-style-type: none"> Should only be administered by a health care provider 	SC: <ul style="list-style-type: none"> 100 mg/mL PFS 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light until the time of use Do not freeze Do not shake 	Upper respiratory infections, injection site reactions and diarrhea
Inflectra (infliximab-dyyb)	Celltrion/Pfizer	<ul style="list-style-type: none"> AS CD (6+) PsO PsA RA UC (6+) 	IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg lyophilized in a powder single dose vial for reconstitution 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) 	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Infliximab (unbranded version of Remicade)	Janssen	<ul style="list-style-type: none"> AS CD (6+) PsO PsA RA UC (6+) 	IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg lyophilized in a powder single dose vial for reconstitution 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) 	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Kevzara (sarilumab)	Sanofi	<ul style="list-style-type: none"> RA PJIA (> 63 kg) PMR 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 150 mg/1.14 mL PFS 200 mg/1.14 mL PFS 150 mg/1.14 mL PEN 200 mg/1.14 mL PEN 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) in original carton to protect from light Do not freeze Do not shake 	RA: Neutropenia, increased ALT, injection site erythema, upper respiratory infections and urinary tract infections PMR: neutropenia, leukopenia and injection site pruritus

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Litfulo (ritlecitinib)	Pfizer	<ul style="list-style-type: none"> • AA (12+) 	Oral	Oral: <ul style="list-style-type: none"> • 50 mg capsules 	<ul style="list-style-type: none"> • Store at 68°F to 77°F (20°C to 25°C) • Store in original package 	Headache, diarrhea, acne, rash, urticaria, folliculitis, pyrexia, atopic dermatitis, dizziness, blood creatine phosphokinase increased, herpes zoster, red blood cell count decreased and stomatitis
Nemluvio (nemolizumab-ilto)	Galderma	<ul style="list-style-type: none"> • AD (12+) • PN 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 30 mg/0.49 mL PEN 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) in original carton to protect from light • Do not freeze 	Headache (including migraine), arthralgia, urticaria, and myalgia.

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Olumiant (baricitinib)	Eli Lilly	<ul style="list-style-type: none"> • AA • RA • Hospitalized adults who have COVID-19 	Oral	Oral: <ul style="list-style-type: none"> • 1 mg tablets • 2 mg tablets • 4 mg tablets 	<ul style="list-style-type: none"> • Store at 68°F to 77°F (20°C to 25°C) 	RA: upper respiratory tract infections, nausea, herpes simplex and herpes zoster COVID-19: increases of liver enzymes, thrombocytosis, creatine phosphokinase increases, neutropenia, deep vein thrombosis, pulmonary embolism and urinary tract infection AA: upper respiratory tract infections, headache, acne, hyperlipidemia, creatine phosphokinase increase, urinary tract infection, liver enzyme elevations, folliculitis, fatigue, lower respiratory tract infections, nausea, genital Candida infections, anemia, neutropenia, abdominal pain, herpes zoster and weight increase
Omvo (mirikizumab-mrkz)	Eli Lilly	<ul style="list-style-type: none"> • UC 	IV infusion: <ul style="list-style-type: none"> • Infused by a health care professional SC injection: <ul style="list-style-type: none"> • Self-administered 	IV: <ul style="list-style-type: none"> • 300 mg/15 mL single dose vial SC: <ul style="list-style-type: none"> • 100 mg/mL PEN 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in original container to protect from light • Do not shake • Do not freeze 	Induction: upper respiratory tract infections and arthralgia. Maintenance: upper respiratory tract infections, injection site reactions, arthralgia, rash, headache and herpes viral infection

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Orencia (abatacept)	Bristol Myers Squibb	IV: <ul style="list-style-type: none"> PJIA (6+) Prophylaxis of aGVHD (2+) PsA RA SC: <ul style="list-style-type: none"> PJIA (2+) PsA (2+) RA 	IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional SC injection: <ul style="list-style-type: none"> Self-administered 	IV: <ul style="list-style-type: none"> 250 mg lyophilized powder in a single dose vial for reconstitution SC: <ul style="list-style-type: none"> 50 mg/0.4 mL PFS 87.5 mg/0.7 mL PFS 125 mg/mL PFS 125 mg/mL PEN 	IV: <ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Store in original container Protect from light SC: <ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Store in original container Protect from light Do not freeze 	IV: <p>RA: headache, upper respiratory tract infection, nasopharyngitis and nausea</p> <p>aGVHD: anemia, hypertension, CMV reactivation/CMV infection, pyrexia, pneumonia, epistaxis, CD4 lymphocytes decreased, hypermagnesemia and acute kidney injury</p> <p>SC: Headache, upper respiratory tract infection, nasopharyngitis and nausea</p>
Otezla (apremilast)	Amgen	<ul style="list-style-type: none"> Oral ulcers associated with Behçet's Disease PsA PsO (6+) 	Oral	Oral: <ul style="list-style-type: none"> 28 day Starter Pack: 10 mg, 20 mg, and 30 mg tablets 20 mg tablets 30 mg tablets 	<ul style="list-style-type: none"> Store below 86°F (30°C) 	<p>PsA: diarrhea, nausea and headache</p> <p>PsO: diarrhea, nausea, upper respiratory tract infection and headache, including tension headache</p> <p>Behçet's Disease: diarrhea, nausea, headache and upper respiratory tract infection</p>

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Otrexup (methotrexate)	Antares Pharma/Otter Pharmaceuticals	<ul style="list-style-type: none"> • PJIA • PsO • RA 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 10 mg/0.4 mL PEN • 12.5 mg/0.4 mL PEN • 15 mg/0.4 mL PEN • 17.5 mg/0.4 mL PEN • 20 mg/0.4 mL PEN • 22.5 mg/0.4 mL PEN • 25 mg/0.4mL PEN 	<ul style="list-style-type: none"> • Store at controlled room temperature, 77°F (25°C); excursions permitted to 59°F to 86°F (15°C to 30°C) • Protect from light 	Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and “burning of skin lesions”

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rasuvo (methotrexate)	Medexus	<ul style="list-style-type: none"> • PJIA • PsO • RA 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 7.5 mg/0.15 mL PEN • 10 mg/0.20 mL PEN • 12.5 mg/0.25 mL PEN • 15 mg/0.30 mL PEN • 17.5 mg/0.35 mL PEN • 20 mg/0.40 mL PEN • 22.5mg/0.45mL PEN • 25 mg/0.50 mL PEN • 27.5 mg/0.55 mL PEN • 30 mg/0.60 mL 	<ul style="list-style-type: none"> • Store at controlled room temperature, 77°F (25°C) • Protect from light 	Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and “burning of skin lesions”
Remicade (infliximab)	Janssen	<ul style="list-style-type: none"> • AS • CD (6+) • PsO • PsA • RA • UC (6+) 	IV infusion: <ul style="list-style-type: none"> • Prepared and infused by a health care professional 	IV: <ul style="list-style-type: none"> • 100 mg lyophilized powder in a single dose vial for reconstitution 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) 	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Renflexis (infliximab-abda)	Samsung Bioepis/ Organon	<ul style="list-style-type: none"> • AS • CD (6+) • PsO • PsA • RA • UC (6+) 	IV infusion: <ul style="list-style-type: none"> • Prepared and infused by a health care professional 	IV: <ul style="list-style-type: none"> • 100 mg lyophilized powder in a single dose vial for reconstitution 	<ul style="list-style-type: none"> • Refrigerate at 36°F to 46°F (2°C to 8°C) 	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Riabni (rituximab-arrx)	Amgen	<ul style="list-style-type: none"> • RA 	IV infusion only: <ul style="list-style-type: none"> • Administer only by a health care professional • Do not administer by IV push or bolus 	IV: <ul style="list-style-type: none"> • 100 mg/10 mL single dose vial • 500 mg/50 mL single dose vial 	<ul style="list-style-type: none"> • Refrigerate vials at 36°F to 46°F (2°C to 8°C) • Do not freeze or shake • Protect from direct sunlight 	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rinvoq (upadacitinib)	AbbVie	<ul style="list-style-type: none"> • AD (12+) • AS • CD • Nr-axSpA • PsA • RA • UC 	Oral	<p>Oral:</p> <ul style="list-style-type: none"> • Extended-release tablets: 15 mg, 30 mg, 45 mg <p>Oral Solution (LQ):</p> <ul style="list-style-type: none"> • 1 mg/1 mL only approved for PJIA (2+) and PsA (2+) 	<ul style="list-style-type: none"> • Store at 36°F to 77°F (2°C to 25°C) • Store in the original bottle to protect from moisture 	<p>RA/PsA/AS/nr-axSpA: upper respiratory tract infections, herpes zoster, herpes simplex, bronchitis, nausea, cough, pyrexia, acne and headache</p> <p>AD: upper respiratory tract infections, acne, herpes simplex, headache, blood creatine phosphokinase increased, cough, hypersensitivity, folliculitis, nausea, abdominal pain, pyrexia, increased weight, herpes zoster, influenza, fatigue, neutropenia, myalgia and influenza like illness</p> <p>UC: upper respiratory tract infections, increased blood creatine phosphokinase, acne, neutropenia, elevated liver enzymes and rash</p> <p>CD: upper respiratory tract infections, anemia, pyrexia, acne, herpes zoster and headache</p>

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rituxan (rituximab)	Genentech	<ul style="list-style-type: none"> RA 	IV infusion only: <ul style="list-style-type: none"> Administer only by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg/10 mL single dose vial 500 mg/50 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate vials at 36°F to 46°F (2°C to 8°C) Do not freeze or shake Protect from direct sunlight 	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Ruxience (rituximab-pvvr)	Pfizer	<ul style="list-style-type: none"> RA 	IV infusion only: <ul style="list-style-type: none"> Administer only by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg/10 mL single dose vial 500 mg/50 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate vials at 36°F to 46°F (2°C to 8°C) Do not freeze or shake Protect from direct sunlight 	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Saphnelo (anifrolumab-fnia)	AstraZeneca	<ul style="list-style-type: none"> SLE 	IV Infusion: <ul style="list-style-type: none"> Infused by a health care professional 	IV: <ul style="list-style-type: none"> 300 mg/2 mL single dose vial 	<ul style="list-style-type: none"> Store in a refrigerator at 36°F to 46°F (2°C to 8°C) Store in the original carton to protect from light Do not freeze Do not shake 	Nasopharyngitis, upper respiratory tract infections, bronchitis, infusion related reactions, herpes zoster and cough

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Siliq (brodalumab)	Valeant/Ortho Dermatologics	<ul style="list-style-type: none"> PsO 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 210 mg/1.5 mL PFS 	<ul style="list-style-type: none"> Store refrigerated at 36°F to 46°F (2°C to 8°C) Store in the original carton to protect from light Do not freeze Do not shake 	Arthralgia, headache, fatigue, diarrhea, oropharyngeal pain, nausea, myalgia, injection site reactions, influenza, neutropenia and tinea infections
Simlandi (adalimumab-ryvk)	Alvotect/Teva	<ul style="list-style-type: none"> AS CD (6+) HS PJIA (2+) PsO PsA RA UC UV 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 40 mg/0.4 mL PEN 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Protect from light Store in original carton 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Simponi ARIA (golimumab)	Janssen	<ul style="list-style-type: none"> AS PJIA (2+) PsA (2+) RA 	IV infusion only: <ul style="list-style-type: none"> Infused by a health care professional 	IV: <ul style="list-style-type: none"> 50 mg/4 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Do not shake Store in original carton to protect from light 	Upper respiratory tract infection, alanine aminotransferase increased, viral infection, aspartate aminotransferase increased, neutrophil count decreased, bronchitis, hypertension and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Skyrizi (risankizumab-rzaa)	AbbVie	IV/OBI: <ul style="list-style-type: none"> • CD • UC SC: <ul style="list-style-type: none"> • PsA • PsO 	IV infusion: <ul style="list-style-type: none"> • Infused by a health care professional SC injection: <ul style="list-style-type: none"> • Self-administered 	IV: <ul style="list-style-type: none"> • 600 mg/10 mL single dose vial SC: <ul style="list-style-type: none"> • 360 mg/2.4 mL OBI • 150 mg/mL PEN • 150 mg/mL PFS 	<ul style="list-style-type: none"> • Store in a refrigerator at 36°F to 46° F (2°C to 8°C) • Do not freeze • Do not shake • Store in the original cartons to protect from light 	PsO and PsA: upper respiratory infections, headache, fatigue, injection site reactions and tinea infections CD: Induction: upper respiratory infections, headache, and arthralgia. Maintenance: arthralgia, abdominal pain, injection site reactions, anemia, pyrexia, back pain, arthropathy and urinary tract infection
Sotyktu (deucravacitinib)	Bristol Meyers Squibb	<ul style="list-style-type: none"> • PsO 	Oral	Oral: <ul style="list-style-type: none"> • 6 mg tablets 	Store at 68°F to 77°F (20°C to 25°C)	Upper respiratory infections, blood creatine phosphokinase increased, herpes simplex, mouth ulcers, folliculitis and acne

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Stelara (ustekinumab)	Janssen	<ul style="list-style-type: none"> • CD • PsA (6+) • PsO (6+) • UC 	IV infusion: <ul style="list-style-type: none"> • Prepared and infused by a health care professional SC injection: <ul style="list-style-type: none"> • Self-administered 	IV: <ul style="list-style-type: none"> • 130 mg/26 mL single dose vial SC: <ul style="list-style-type: none"> • 45 mg/0.5 mL PFS • 90 mg/mL PFS • 45 mg/0.5 mL single-dose vial 	<ul style="list-style-type: none"> • Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light 	PsO: nasopharyngitis, upper respiratory tract infection, headache and fatigue CD: Induction: vomiting. Maintenance: nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic infection, bronchitis, pruritus, urinary tract UC: Induction: nasopharyngitis. Maintenance: nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea
Taltz (ixekizumab)	Eli Lilly	<ul style="list-style-type: none"> • AS • Nr-axSpA • PsA • PsO (6+) 	SC injection: <ul style="list-style-type: none"> • Self-administered 	SC: <ul style="list-style-type: none"> • 20 mg/0.25 mL PFS • 40 mg/0.5 mL PFS • 80 mg/mL PEN • 80 mg/mL PFS 	<ul style="list-style-type: none"> • Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Do not shake 	Injection site reactions, upper respiratory tract infections, nausea and tinea infections

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Tremfya (guselkumab)	Janssen	<ul style="list-style-type: none"> PsA PsO UC 	SC injection: <ul style="list-style-type: none"> Self-administered IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional 	SC: <ul style="list-style-type: none"> 100 mg/mL PFS 100 mg/mL PEN 200 mg/2 mL PFS 200 mg/2 mL PEN IV: <ul style="list-style-type: none"> 200 mg/20 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Store in original carton to protect from light Do not freeze Do not shake 	Upper respiratory infections, headache, injection site reactions, arthralgia, bronchitis, diarrhea, gastroenteritis, tinea infections and herpes simplex infections
Truxima (rituximab-abbs)	Celltrion/Teva	<ul style="list-style-type: none"> RA 	IV infusion only: <ul style="list-style-type: none"> Administer only by a health care professional 	IV: <ul style="list-style-type: none"> 100 mg/10 mL single dose vial 500 mg/50 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate vials at 36°F to 46°F (2°C to 8°C) Do not freeze or shake Protect from direct sunlight 	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Tyenne (tocilizumab-aazg)	Fresenius Kabi	<ul style="list-style-type: none"> GCA PJIA (2+) RA SJIA (2+) 	SC Injection: <ul style="list-style-type: none"> Self-administered IV infusion: <ul style="list-style-type: none"> Prepared and infused by a health care professional 	SC: <ul style="list-style-type: none"> 162 mg/0.9 mL PFS 162 mg/0.9 mL PEN IV: <ul style="list-style-type: none"> 80 mg/4 mL single dose vial 200 mg/10 mL single dose vial 400 mg/20 mL single dose vial 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Store in original packages Protect from light Keep syringes dry 	Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Velsipity (etrasimod)	Pfizer	<ul style="list-style-type: none"> UC 	Oral	Oral: <ul style="list-style-type: none"> 2 mg tablet 	<ul style="list-style-type: none"> Store at room temperature at 68°F to 77°F (20°C to 25°C) 	Headache, elevated liver tests and dizziness
Xeljanz/Xeljanz XR (tofacitinib)	Pfizer	<ul style="list-style-type: none"> AS PJIA (2+) PsA RA UC 	Oral	Oral: <ul style="list-style-type: none"> Tablets: <ul style="list-style-type: none"> 5 mg 10 mg XR tablets: <ul style="list-style-type: none"> 11 mg 22 mg Oral Solution: <ul style="list-style-type: none"> 1 mg/mL 	<ul style="list-style-type: none"> Store at room temperature at 68°F to 77°F (20°C to 25°C) Do not repackage 	RA, PsA, AS and PJIA: upper respiratory tract infection, nasopharyngitis, diarrhea and headache UC: nasopharyngitis, elevated cholesterol levels, headache, upper respiratory tract infection, increased blood creatine phosphokinase, rash, diarrhea and herpes zoster
Yuflyma (adalimumab-aaty)	Celltrion	<ul style="list-style-type: none"> AS CD (6+) HS PJIA (2+) PsO PsA RA UC UV 	SC injection: <ul style="list-style-type: none"> Self-administered 	SC: <ul style="list-style-type: none"> 20 mg/0.2 mL PFS 40 mg/0.4 mL PFS 40 mg/0.4 mL PEN 80 mg/0.8 mL PEN 	<ul style="list-style-type: none"> Refrigerate at 36°F to 46°F (2°C to 8°C) Do not freeze Protect from light 	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Zeposia (ozanimod)	Bristol Myers Squibb	• UC	Oral	Oral: • 7 Day starter pack: – 0.23 mg capsules – 0.46 mg capsules • 0.92 mg capsules	• Store at 68°F to 77°F (20°C to 25°C)	Liver test increased, upper respiratory infection and headache



Visit cvs.co/ai-list or scan the QR code to view electronically.
For easy access, you can bookmark this page.

FDA (U.S. Food and Drug Administration). SC (Subcutaneous). IV (Intravenous). PFS (Prefilled syringe). OBI (On-body injector).

AA (Alopecia areata). AD (Atopic dermatitis). aGVHD (Acute graft versus host disease). AOSD (Adult-onset Still's Disease). AS (Ankylosing spondylitis). CAPS (Cryopyrin-associated periodic syndrome). CRS (Cytokine release syndrome). COPD (Chronic obstructive pulmonary disease). CRSwNP (Chronic rhinosinusitis with nasal polyposis). DIRA (Deficiency of IL-1-receptor antagonist). EoE (Eosinophilic esophagitis). ERA (Enthesitis-related arthritis). FCAS (Familial cold autoinflammatory syndrome). FMF (Familial Mediterranean fever). GCA (Giant cell arteritis). HIDS/MKD (Hyperimmunoglobulin D syndrome / mevalonate kinase deficiency). HS (Hidradenitis Suppurativa). JIA (Juvenile idiopathic arthritis). LN (Lupus Nephritis). MWS (Muckle-Wells syndrome). NOMID (Neonatal-onset multisystem inflammatory disease). Nr-axSpA (Non-radiographic axial spondyloarthritis). PJIA (Polyarticular Juvenile idiopathic arthritis). PMR (Polymyalgia rheumatica). PN (Prurigo nodularis). PsA (Psoriatic arthritis). PsO (Plaque psoriasis). RA (Rheumatoid arthritis). RP (Recurrent pericarditis). SJIA (Systemic Juvenile idiopathic arthritis). SLE (Systemic lupus erythematosus). SSc-ILD (Systemic sclerosis - Interstitial lung disease). TRAPS (Tumor necrosis factor receptor-associated periodic syndrome). UC (Ulcerative colitis). UV (Uveitis).

The information contained in this communication is provided in summary form. It is not intended for use as the sole basis of clinical treatment, as a substitute for reading the original research, nor as a substitute for the knowledge, skill and judgment of the medical provider. This list is subject to change. CVS Specialty assumes no liability whatsoever for the information provided or for any diagnosis or treatment made as a result of this information. The information noted in this brochure was taken from the manufacturers' prescribing information. It does not contain all information needed to use these products safely and effectively. See the manufacturers' package insert for full prescribing information. CVS Specialty does not provide medical advice and does not recommend a particular treatment or brand. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Specialty.