



Visit [cvs.co/ai-list](https://cvs.co/ai-list) or scan the QR code to view electronically.  
For easy access, you can bookmark this page.

## Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name   | Manufacturer   | FDA-Approved Indication(s)  | Route of Administration   | Supply Method   | Storage  | Select Common Adverse Reactions  |
|---|----------------|---|---|---|--|--|
| <b>Actemra</b><br>(tocilizumab)                         | Genentech      | <ul style="list-style-type: none"> <li>• CRS (2+)</li> <li>• GCA</li> <li>• PJIA (2+)</li> <li>• RA</li> <li>• SJIA (2+)</li> <li>• SSc-ILD</li> <li>• Hospitalized adults who have COVID-19</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 162 mg/0.9 mL PFS</li> <li>• 162 mg/0.9 mL PEN</li> </ul> <b>IV:</b> <ul style="list-style-type: none"> <li>• 80 mg/4 mL</li> <li>• 200 mg/10 mL</li> <li>• 400 mg/20 mL</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Store in original packages</li> <li>• Protect from light</li> <li>• Keep syringes dry</li> </ul> | Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions |
| <b>Adalimumab-aacf</b><br>(unbranded version of Idacio) | Fresenius Kabi | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS (12+)</li> <li>• PJIA (2+)</li> <li>• PsA</li> <li>• PsO</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul>              | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>  | <b>SC:</b> <ul style="list-style-type: none"> <li>• 40 mg/0.8 mL PEN</li> </ul>   | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Store in original packages</li> <li>• Protect from light</li> </ul>                              | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash   |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name  | Manufacturer  | FDA-Approved Indication(s)  | Route of Administration  | Supply Method   | Storage   | Select Common Adverse Reactions  |
|--|---------------|---|--|---|---|--|
| <b>Adalimumab-aaty</b><br>(unbranded version of Yuflyma) | Celltrion USA | <b>SC injection:</b> <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>HS (12+)</li> <li>PJIA (2+)</li> <li>PsA</li> <li>PsO</li> <li>RA</li> <li>UC</li> <li>UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>20 mg/0.2 mL PFS</li> <li>40 mg/0.4 mL PFS</li> <li>40 mg/0.4 mL PEN</li> <li>80 mg/0.8 mL PEN</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Store in original packages</li> <li>Protect from light</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |
| <b>Adalimumab-adaz</b><br>(unbranded version of Hyrimoz) | Sandoz        | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>HS</li> <li>PJIA (2+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC</li> <li>UV</li> </ul>                            | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>40 mg/0.4 mL PEN</li> <li>40 mg/0.4 mL PFS</li> </ul>   | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Store in original packages</li> <li>Protect from light</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name  | Manufacturer | FDA-Approved Indication(s)   | Route of Administration  | Supply Method   | Storage  | Select Common Adverse Reactions   |
|--|--------------|--|--|---|--|---|
| <b>Adalimumab-fkjp</b><br>(unbranded version of Hulio) | Mylan        | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>HS</li> <li>PJIA (2+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC</li> <li>UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>40 mg/0.8 mL PEN</li> <li>40 mg/0.8 mL PFS</li> <li>20 mg/0.4 mL PFS</li> </ul>                           | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Protect from light</li> <li>Store in original package</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash  |
| <b>Adbry</b><br>(tralokinumab-ldrm)                    | LEO Pharma   | <ul style="list-style-type: none"> <li>AD (12+)</li> </ul>   | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>150 mg/mL PFS</li> <li>300 mg/2 mL PEN</li> </ul>   | <ul style="list-style-type: none"> <li>Store refrigerated at 36°F to 46°F (2°C to 8°C)</li> <li>Store in the original carton to protect from light</li> </ul>                    | Upper respiratory tract infections, conjunctivitis, injection site reactions and eosinophilia |
| <b>Amjevita</b><br>(adalimumab-atto)                   | Amgen        | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>HS</li> <li>PJIA (2+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC</li> <li>UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>40 mg/0.8 mL PEN</li> <li>40 mg/0.8 mL PFS</li> <li>20 mg/0.4 mL PFS</li> <li>10 mg/0.2 mL PFS</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Protect from light</li> <li>Store in original package</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                          | Manufacturer            | FDA-Approved Indication(s)  | Route of Administration  | Supply Method  | Storage  | Select Common Adverse Reactions  |
|------------------------------------|-------------------------|---|--|--|--|--|
| <b>Arcalyst</b><br>(rilonacept)    | Kiniksa Pharmaceuticals | <ul style="list-style-type: none"> <li>• CAPS (12+)</li> <li>• FCAS (12+)</li> <li>• MWS (12+)</li> <li>• DIRA (<math>\geq 10</math> kg)</li> <li>• RP (12+)</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>                                 | <b>SC:</b> <ul style="list-style-type: none"> <li>• 220 mg of lyophilized powder in a single dose vial for reconstitution</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Store in original carton</li> <li>• Protect from light</li> <li>• Do not use beyond the date stamped on the label</li> <li>• After reconstitution, may be kept at room temperature, protected from light</li> </ul> | Injection-site reactions and upper respiratory tract infections  |
| <b>Avsola</b><br>(infliximab-axxq) | Amgen                   | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC (6+)</li> </ul>                                | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 100 mg lyophilized powder in a single dose vial for reconstitution</li> </ul>    | <ul style="list-style-type: none"> <li>• Store unopened Avsola vials in a refrigerator at 36°F to 46°F (2°C to 8°C)</li> <li>• Protect from light</li> </ul>   | Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                            | Manufacturer    | FDA-Approved Indication(s)  | Route of Administration  | Supply Method  | Storage   | Select Common Adverse Reactions  |
|--------------------------------------|-----------------|---|--|--|---|--|
| <b>Benlysta</b><br>(belimumab)       | GlaxoSmithKline | <ul style="list-style-type: none"> <li>• SLE (5+)</li> <li>• LN (5+)</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 120 mg lyophilized powder in a single dose vial for reconstitution</li> <li>• 400 mg lyophilized powder in a single dose vial for reconstitution</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>• 200 mg/mL PEN</li> <li>• 200 mg/mL PFS</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Store in original packages</li> <li>• Protect from light</li> <li>• Avoid exposure to heat</li> </ul> | Nausea, diarrhea, pyrexia, nasopharyngitis, bronchitis, pharyngitis, insomnia, pain in extremity, depression and migraine  |
| <b>Bimzelx</b><br>(bimekizumab-bkzx) | UCB             | <ul style="list-style-type: none"> <li>• PsO</li> </ul>                         | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>   | <b>SC:</b> <ul style="list-style-type: none"> <li>• 160mg/mL PEN</li> <li>• 160mg/mL PFS</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Do not shake</li> <li>• Store in original carton to protect from light</li> </ul>                     | Upper respiratory tract infections, oral candidiasis, headache, injection site reactions, tinea infections, gastroenteritis, Herpes simplex infections, acne, folliculitis, other candida infections and fatigue |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                             | Manufacturer | FDA-Approved Indication(s)  | Route of Administration   | Supply Method   | Storage  | Select Common Adverse Reactions  |
|---------------------------------------|--------------|---|---|---|--|--|
| <b>Cibinqo</b><br>(abrocitinib)       | Pfizer       | <ul style="list-style-type: none"> <li>AD (12+)</li> </ul>  | Oral  | <b>Oral:</b> <ul style="list-style-type: none"> <li>50 mg tablets</li> <li>100 mg tablets</li> <li>200 mg tablets</li> </ul>  | <ul style="list-style-type: none"> <li>Store at 68°F to 77°F (20°C to 25°C)</li> <li>Keep in original package</li> </ul>   | Nasopharyngitis, nausea, headache, herpes simplex, increased blood creatinine phosphokinase, dizziness, urinary tract infection, fatigue, acne, vomiting, oropharyngeal pain, influenza, gastroenteritis, impetigo, hypertension, contact dermatitis, upper abdominal pain, abdominal discomfort, herpes zoster and thrombocytopenia |
| <b>Cimzia</b><br>(certolizumab pegol) | UCB          | <ul style="list-style-type: none"> <li>AS</li> <li>CD</li> <li>Nr-axSpA</li> <li>PsO</li> <li>PsA</li> <li>RA</li> </ul>                            | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered (PFS only)</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>200 mg/mL PFS</li> <li>200 mg lyophilized powder in a single-dose vial (for health care professional use only)</li> </ul>   | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Protect from light</li> </ul>  | Upper respiratory tract infection, rash and urinary tract infection  |
| <b>Cosentyx</b><br>(secukinumab)      | Novartis     | <b>SC:</b> <ul style="list-style-type: none"> <li>AS</li> <li>ERA (4+)</li> <li>HS</li> <li>Nr-axSpA</li> <li>PsO (6+)</li> <li>PsA (2+)</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul>            | <b>SC:</b> <ul style="list-style-type: none"> <li>75 mg/0.5 mL PFS</li> <li>150 mg/mL PEN</li> <li>150 mg/mL PFS</li> <li>300 mg/2 mL PEN</li> <li>300 mg/2 mL PFS</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Keep the product in the original carton</li> <li>Protect from light</li> <li>Do not shake to avoid foaming</li> </ul> | Nasopharyngitis, diarrhea and upper respiratory tract infection  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                      | Manufacturer | FDA-Approved Indication(s)  | Route of Administration  | Supply Method   | Storage  | Select Common Adverse Reactions   |
|--------------------------------|--------------|---|--|---|--|---|
| <b>Dupixent</b><br>(dupilumab) | Sanofi       | <ul style="list-style-type: none"> <li>• AD (6 mo+)</li> <li>• Asthma (6+)</li> <li>• CRSwNP</li> <li>• EoE (1+ and ≥ 15 kg)</li> <li>• PN</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 100 mg/0.67 mL PFS</li> <li>• 200 mg/1.14 mL PEN</li> <li>• 200 mg/1.14 mL PFS</li> <li>• 300 mg/2 mL PEN</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light</li> <li>• Do not freeze</li> </ul>   | Injection site reactions, conjunctivitis, blepharitis, oral herpes, keratitis, eye pruritus, other herpes simplex virus infection and dry eye |
| <b>Enbrel</b><br>(etanercept)  | Amgen        | <ul style="list-style-type: none"> <li>• AS</li> <li>• JPsA (2+)</li> <li>• PsO (4+)</li> <li>• PsA (2+)</li> <li>• RA</li> </ul>                     | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 25 mg/0.5 mL single dose vial</li> <li>• 25 mg lyophilized powder in a multiple dose vial for reconstitution</li> <li>• 25 mg/0.5 mL PFS</li> <li>• 50 mg/mL PFS</li> <li>• 50 mg/mL PEN</li> <li>• 50 mg/mL Enbrel Mini single-dose prefilled cartridge</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Do not shake</li> <li>• Store in the original carton to protect from light or physical damage</li> </ul> | Infections and injection site reactions   |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                           | Manufacturer                | FDA-Approved Indication(s)   | Route of Administration   | Supply Method   | Storage  | Select Common Adverse Reactions  |
|-------------------------------------|-----------------------------|--|---|---|--|--|
| <b>Entyvio</b><br>(vedolizumab)     | Takeda                      | <ul style="list-style-type: none"> <li>• CD</li> <li>• UC</li> </ul>   | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Administered by a healthcare provider</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 300 mg single dose vial</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>• 108 mg/0.68 mL PEN</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate unopened vials, PEN, and PFS at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Retain in original package to protect from light</li> </ul> | <p>Nasopharyngitis, headache, arthralgia, nausea, pyrexia, upper respiratory tract infection, fatigue, cough, bronchitis, influenza, back pain, rash, pruritus, sinusitis, oropharyngeal pain and pain in extremities.</p> <p>In addition, injection site reactions are associated with the SC injection only.</p> |
| <b>Hadlima</b><br>(adalimumab-bwwd) | Samsung Bioepis/<br>Organon | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>  | <b>SC:</b> <ul style="list-style-type: none"> <li>• 40 mg/0.8 mL PFS</li> <li>• 40 mg/0.4 mL PFS</li> <li>• 40 mg/0.8 mL PEN</li> <li>• 40 mg/0.4 mL PEN</li> </ul>         | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> <li>• Store in original container</li> </ul>                     | <p>Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash</p>  |



# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                         | Manufacturer        | FDA-Approved Indication(s)   | Route of Administration  | Supply Method   | Storage   | Select Common Adverse Reactions  |
|-----------------------------------|---------------------|--|--|---|---|--|
| <b>Hulio</b><br>(adalimumab-fkjp) | Mylan               | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul>                 | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 40 mg/0.8 mL PEN</li> <li>• 40 mg/0.8 mL PFS</li> <li>• 20 mg/0.4 mL PFS</li> </ul>   | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> <li>• Store in original carton</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |
| <b>Humira</b><br>(adalimumab)     | AbbVie/<br>Cordavis | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS (12+)</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC (5+)</li> <li>• UV (2+)</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 80 mg/0.8 mL PEN</li> <li>• 40 mg/0.4 mL PEN</li> <li>• 40 mg/0.4 mL PFS</li> <li>• 20 mg/0.2 mL PFS</li> <li>• 10 mg/0.1 mL PFS</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> <li>• Store in original carton</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                           | Manufacturer        | FDA-Approved Indication(s)   | Route of Administration  | Supply Method  | Storage   | Select Common Adverse Reactions  |
|-------------------------------------|---------------------|--|--|--|---|--|
| <b>Hyrimoz</b><br>(adalimumab-adaz) | Cordavis/<br>Sandoz | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 80 mg/0.8 mL PEN</li> <li>• 40 mg/0.4 mL PEN</li> <li>• 40 mg/0.8 mL PEN</li> <li>• 80 mg/0.8 mL PFS</li> <li>• 40 mg/0.4 mL PFS</li> <li>• 40 mg/0.8 mL PFS</li> <li>• 20 mg/0.2 mL PFS</li> <li>• 10 mg/ 0.1 mL PFS</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> <li>• Store in original carton</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |
| <b>Idacio</b><br>(adalimumab-aacf)  | Fresenius Kabi      | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 40 mg/0.8 mL PEN</li> <li>• 40 mg/0.8 mL PFS</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> </ul>                                     | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                             | Manufacturer | FDA-Approved Indication(s)   | Route of Administration  | Supply Method   | Storage  | Select Common Adverse Reactions  |
|---------------------------------------|--------------|--|--|---|--|--|
| <b>Ilaris</b><br>(canakinumab)        | Novartis     | <ul style="list-style-type: none"> <li>• CAPS (4+)</li> <li>• FCAS (4+)</li> <li>• FMF</li> <li>• Gout flares</li> <li>• HIDS/MKD</li> <li>• MWS (4+)</li> <li>• TRAPS</li> <li>• Active Still's Disease/AOSD (2+)</li> <li>• SJIA (2+)</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Should only be administered by a health care professional</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 150 mg/mL single dose vial</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Store in the original carton to protect from light</li> </ul>                              | <b>CAPS:</b> Nasopharyngitis, diarrhea, influenza, rhinitis, nausea, headache, bronchitis, gastroenteritis, pharyngitis, weight increased, musculoskeletal pain and vertigo<br><b>TRAPS, HIDS/MKD, and FMF:</b> injection-site reactions and nasopharyngitis<br><b>Still's Disease:</b> Nasopharyngitis and upper respiratory tract infections, abdominal pain and injection-site reactions<br><b>Gout Flares:</b> Nasopharyngitis, upper respiratory tract infections, urinary tract infections, hypertriglyceridemia and back pain |
| <b>Ilumya</b><br>(tildrakizumab-asmn) | Sun Pharma   | PsO  | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Should only be administered by a health care provider</li> </ul>     | <b>SC:</b> <ul style="list-style-type: none"> <li>• 100 mg/mL PFS</li> </ul>              | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light until the time of use</li> <li>• Do not freeze</li> <li>• Do not shake</li> </ul> | Upper respiratory infections, injection site reactions and diarrhea  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name  | Manufacturer     | FDA-Approved Indication(s)   | Route of Administration  | Supply Method   | Storage  | Select Common Adverse Reactions  |
|--|------------------|--|--|---|--|--|
| <b>Inflectra</b><br>(infliximab-dyyb)                | Celltrion/Pfizer | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC (6+)</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>Prepared and infused by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>100 mg lyophilized in a powder single dose vial for reconstitution</li> </ul>                                     | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> </ul>   | Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain   |
| <b>Infliximab</b><br>(unbranded version of Remicade) | Janssen          | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC (6+)</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>Prepared and infused by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>100 mg lyophilized in a powder single dose vial for reconstitution</li> </ul>                                     | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> </ul>   | Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain   |
| <b>Kevzara</b><br>(sarilumab)                        | Sanofi           | <ul style="list-style-type: none"> <li>RA</li> <li>PJIA (&gt; 63 kg)</li> <li>PMR</li> </ul>                                 | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul>                                 | <b>SC:</b> <ul style="list-style-type: none"> <li>150 mg/1.14 mL PFS</li> <li>200 mg/1.14 mL PFS</li> <li>150 mg/1.14 mL PEN</li> <li>200 mg/1.14 mL PEN</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C) in original carton to protect from light</li> <li>Do not freeze</li> <li>Do not shake</li> </ul> | <b>RA:</b> Neutropenia, increased ALT, injection site erythema, upper respiratory infections and urinary tract infections<br><br><b>PMR:</b> neutropenia, leukopenia and injection site pruritus |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                        | Manufacturer | FDA-Approved Indication(s)  | Route of Administration | Supply Method   | Storage   | Select Common Adverse Reactions   |
|----------------------------------|--------------|---|-------------------------|---|---|---|
| <b>Litfulo</b><br>(ritlecitinib) | Pfizer       | <ul style="list-style-type: none"> <li>• AA (12+)</li> </ul>  | Oral                    | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 50 mg capsules</li> </ul>   | <ul style="list-style-type: none"> <li>• Store at 68°F to 77°F (20°C to 25°C)</li> <li>• Store in original package</li> </ul> | Headache, diarrhea, acne, rash, urticaria, folliculitis, pyrexia, atopic dermatitis, dizziness, blood creatine phosphokinase increased, herpes zoster, red blood cell count decreased and stomatitis  |
| <b>Olumiant</b><br>(baricitinib) | Eli Lilly    | <ul style="list-style-type: none"> <li>• AA</li> <li>• RA</li> <li>• Hospitalized adults who have COVID-19</li> </ul> | Oral                    | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 1 mg tablets</li> <li>• 2 mg tablets</li> <li>• 4 mg tablets</li> </ul> | <ul style="list-style-type: none"> <li>• Store at 68°F to 77°F (20°C to 25°C)</li> </ul>                                      | <p><b>RA:</b> upper respiratory tract infections, nausea, herpes simplex and herpes zoster</p> <p><b>COVID-19:</b> increases of liver enzymes, thrombocytosis, creatine phosphokinase increases, neutropenia, deep vein thrombosis, pulmonary embolism and urinary tract infection</p> <p><b>AA:</b> upper respiratory tract infections, headache, acne, hyperlipidemia, creatine phosphokinase increase, urinary tract infection, liver enzyme elevations, folliculitis, fatigue, lower respiratory tract infections, nausea, genital Candida infections, anemia, neutropenia, abdominal pain, herpes zoster and weight increase</p> |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                         | Manufacturer         | FDA-Approved Indication(s)   | Route of Administration  | Supply Method  | Storage   | Select Common Adverse Reactions  |
|-----------------------------------|----------------------|--|--|--|---|--|
| <b>Omvo</b><br>(mirikizumab-mrkz) | Eli Lilly            | <ul style="list-style-type: none"> <li>UC</li> </ul>   | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>Infused by a health care professional</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul>              | <b>IV:</b> <ul style="list-style-type: none"> <li>300 mg/15 mL single dose vial</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>100 mg/mL PEN</li> </ul>   | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Store in original container to protect from light</li> <li>Do not shake</li> <li>Do not freeze</li> </ul>  | <b>Induction:</b> upper respiratory tract infections and arthralgia.<br><b>Maintenance:</b> upper respiratory tract infections, injection site reactions, arthralgia, rash, headache and herpes viral infection  |
| <b>Orencia</b><br>(abatacept)     | Bristol Myers Squibb | <b>IV:</b> <ul style="list-style-type: none"> <li>PJIA (6+)</li> <li>Prophylaxis of aGVHD (2+)</li> <li>PsA</li> <li>RA</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>PJIA (2+)</li> <li>PsA (2+)</li> <li>RA</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>Prepared and infused by a health care professional</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>250 mg lyophilized powder in a single dose vial for reconstitution</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>50 mg/0.4 mL PFS</li> <li>87.5 mg/0.7 mL PFS</li> <li>125 mg/mL PFS</li> <li>125 mg/mL PEN</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Store in original container</li> <li>Protect from light</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Store in original container</li> <li>Protect from light</li> <li>Do not freeze</li> </ul> | <b>IV:</b><br><b>RA:</b> headache, upper respiratory tract infection, nasopharyngitis and nausea<br><b>aGVHD:</b> anemia, hypertension, CMV reactivation/CMV infection, pyrexia, pneumonia, epistaxis, CD4 lymphocytes decreased, hypermagnesemia and acute kidney injury<br><br><b>SC:</b><br>Headache, upper respiratory tract infection, nasopharyngitis and nausea |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                        | Manufacturer                         | FDA-Approved Indication(s)  | Route of Administration  | Supply Method  | Storage  | Select Common Adverse Reactions  |
|----------------------------------|--------------------------------------|---|--|--|--|--|
| <b>Otezla</b><br>(apremilast)    | Amgen                                | <ul style="list-style-type: none"> <li>• Oral ulcers associated with Behçet's Disease</li> <li>• PsA</li> <li>• PsO (6+)</li> </ul> | Oral   | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 28 day Starter Pack: 10 mg, 20 mg, and 30 mg tablets</li> <li>• 20 mg tablets</li> <li>• 30 mg tablets</li> </ul>  | <ul style="list-style-type: none"> <li>• Store below 86°F (30°C)</li> </ul>  | <b>PsA:</b> diarrhea, nausea and headache<br><b>PsO:</b> diarrhea, nausea, upper respiratory tract infection and headache, including tension headache<br><b>Behçet's Disease:</b> diarrhea, nausea, headache and upper respiratory tract infection                             |
| <b>Otrexup</b><br>(methotrexate) | Antares Pharma/Otter Pharmaceuticals | <ul style="list-style-type: none"> <li>• PJIA</li> <li>• PsO</li> <li>• RA</li> </ul>   | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 10 mg/0.4 mL PEN</li> <li>• 12.5 mg/0.4 mL PEN</li> <li>• 15 mg/0.4 mL PEN</li> <li>• 17.5 mg/0.4 mL PEN</li> <li>• 20 mg/0.4 mL PEN</li> <li>• 22.5 mg/0.4 mL PEN</li> <li>• 25 mg/0.4mL PEN</li> </ul> | <ul style="list-style-type: none"> <li>• Store at controlled room temperature, 77°F (25°C); excursions permitted to 59°F to 86°F (15°C to 30°C)</li> <li>• Protect from light</li> </ul> | Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and "burning of skin lesions" |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                       | Manufacturer | FDA-Approved Indication(s)   | Route of Administration  | Supply Method  | Storage   | Select Common Adverse Reactions  |
|---------------------------------|--------------|--|--|--|---|--|
| <b>Rasuvo</b><br>(methotrexate) | Medexus      | <ul style="list-style-type: none"> <li>• PJIA</li> <li>• PsO</li> <li>• RA</li> </ul>  | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>                                 | <b>SC:</b> <ul style="list-style-type: none"> <li>• 7.5 mg/0.15 mL PEN</li> <li>• 10 mg/0.20 mL PEN</li> <li>• 12.5 mg/0.25 mL PEN</li> <li>• 15 mg/0.30 mL PEN</li> <li>• 17.5 mg/0.35 mL PEN</li> <li>• 20 mg/0.40 mL PEN</li> <li>• 22.5mg/0.45mL PEN</li> <li>• 25 mg/0.50 mL PEN</li> <li>• 27.5 mg/0.55 mL PEN</li> <li>• 30 mg/0.60 mL</li> </ul> | <ul style="list-style-type: none"> <li>• Store at controlled room temperature, 77°F (25°C)</li> <li>• Protect from light</li> </ul> | Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and “burning of skin lesions” |
| <b>Remicade</b><br>(infliximab) | Janssen      | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC (6+)</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 100 mg lyophilized powder in a single dose vial for reconstitution</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> </ul>  | Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain   |



# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                             | Manufacturer                | FDA-Approved Indication(s)   | Route of Administration   | Supply Method   | Storage  | Select Common Adverse Reactions   |
|---------------------------------------|-----------------------------|--|---|---|--|---|
| <b>Renflexis</b><br>(infliximab-abda) | Samsung Bioepis/<br>Organon | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC (6+)</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul>  | <b>IV:</b> <ul style="list-style-type: none"> <li>• 100 mg lyophilized powder in a single dose vial for reconstitution</li> </ul>     | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> </ul>   | Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain  |
| <b>Riabni</b><br>(rituximab-arrx)     | Amgen                       | <ul style="list-style-type: none"> <li>• RA</li> </ul>   | <b>IV infusion only:</b> <ul style="list-style-type: none"> <li>• Administer only by a health care professional</li> <li>• Do not administer by IV push or bolus</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 100 mg/10 mL single dose vial</li> <li>• 500 mg/50 mL single dose vial</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate vials at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze or shake</li> <li>• Protect from direct sunlight</li> </ul> | Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events) |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                       | Manufacturer | FDA-Approved Indication(s)  | Route of Administration | Supply Method  | Storage  | Select Common Adverse Reactions  |
|---------------------------------|--------------|---|-------------------------|--|--|--|
| <b>Rinvoq</b><br>(upadacitinib) | AbbVie       | <ul style="list-style-type: none"> <li>• AD (12+)</li> <li>• AS</li> <li>• CD</li> <li>• Nr-axSpA</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> </ul> | Oral                    | <p><b>Oral:</b></p> <ul style="list-style-type: none"> <li>• Extended-release tablets: 15 mg, 30 mg, 45 mg</li> </ul> <p><b>Oral Solution (LQ):</b></p> <ul style="list-style-type: none"> <li>• 1 mg/1 mL only approved for PJIA (2+) and PsA (2+)</li> </ul> | <ul style="list-style-type: none"> <li>• Store at 36°F to 77°F (2°C to 25°C)</li> <li>• Store in the original bottle to protect from moisture</li> </ul> | <p><b>RA/PsA/AS/nr-axSpA:</b> upper respiratory tract infections, herpes zoster, herpes simplex, bronchitis, nausea, cough, pyrexia, acne and headache</p> <p><b>AD:</b> upper respiratory tract infections, acne, herpes simplex, headache, blood creatine phosphokinase increased, cough, hypersensitivity, folliculitis, nausea, abdominal pain, pyrexia, increased weight, herpes zoster, influenza, fatigue, neutropenia, myalgia and influenza like illness</p> <p><b>UC:</b> upper respiratory tract infections, increased blood creatine phosphokinase, acne, neutropenia, elevated liver enzymes and rash</p> <p><b>CD:</b> upper respiratory tract infections, anemia, pyrexia, acne, herpes zoster and headache</p> |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                             | Manufacturer | FDA-Approved Indication(s)                            | Route of Administration  | Supply Method   | Storage   | Select Common Adverse Reactions   |
|---------------------------------------|--------------|---|--|---|---|---|
| <b>Rituxan</b><br>(rituximab)         | Genentech    | <ul style="list-style-type: none"> <li>RA</li> </ul>  | <b>IV infusion only:</b> <ul style="list-style-type: none"> <li>Administer only by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>100 mg/10 mL single dose vial</li> <li>500 mg/50 mL single dose vial</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate vials at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze or shake</li> <li>Protect from direct sunlight</li> </ul>  | Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events) |
| <b>Ruxience</b><br>(rituximab-pvvr)   | Pfizer       | <ul style="list-style-type: none"> <li>RA</li> </ul>  | <b>IV infusion only:</b> <ul style="list-style-type: none"> <li>Administer only by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>100 mg/10 mL single dose vial</li> <li>500 mg/50 mL single dose vial</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate vials at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze or shake</li> <li>Protect from direct sunlight</li> </ul>  | Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events) |
| <b>Saphnelo</b><br>(anifrolumab-fnia) | AstraZeneca  | <ul style="list-style-type: none"> <li>SLE</li> </ul> | <b>IV Infusion:</b> <ul style="list-style-type: none"> <li>Infused by a health care professional</li> </ul>              | <b>IV:</b> <ul style="list-style-type: none"> <li>300 mg/2 mL single dose vial</li> </ul>   | <ul style="list-style-type: none"> <li>Store in a refrigerator at 36°F to 46°F (2°C to 8°C)</li> <li>Store in the original carton to protect from light</li> <li>Do not freeze</li> <li>Do not shake</li> </ul> | Nasopharyngitis, upper respiratory tract infections, bronchitis, infusion related reactions, herpes zoster and cough  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                            | Manufacturer                | FDA-Approved Indication(s)   | Route of Administration  | Supply Method  | Storage  | Select Common Adverse Reactions   |
|--------------------------------------|-----------------------------|--|--|--|--|---|
| <b>Siliq</b><br>(brodalumab)         | Valeant/Ortho Dermatologics | <ul style="list-style-type: none"> <li>PsO</li> </ul>  | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul>                         | <b>SC:</b> <ul style="list-style-type: none"> <li>210 mg/1.5 mL PFS</li> </ul>           | <ul style="list-style-type: none"> <li>Store refrigerated at 36°F to 46°F (2°C to 8°C)</li> <li>Store in the original carton to protect from light</li> <li>Do not freeze</li> <li>Do not shake</li> </ul> | Arthralgia, headache, fatigue, diarrhea, oropharyngeal pain, nausea, myalgia, injection site reactions, influenza, neutropenia and tinea infections   |
| <b>Simlandi</b><br>(adalimumab-ryvk) | Alvotech/Teva               | <ul style="list-style-type: none"> <li>AS</li> <li>CD (6+)</li> <li>HS</li> <li>PJIA (2+)</li> <li>PsO</li> <li>PsA</li> <li>RA</li> <li>UC</li> <li>UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>Self-administered</li> </ul>                         | <b>SC:</b> <ul style="list-style-type: none"> <li>40 mg/0.4 mL PEN</li> </ul>            | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Protect from light</li> <li>Store in original carton</li> </ul>                            | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash  |
| <b>Simponi ARIA</b><br>(golimumab)   | Janssen                     | <ul style="list-style-type: none"> <li>AS</li> <li>PJIA (2+)</li> <li>PsA (2+)</li> <li>RA</li> </ul>  | <b>IV infusion only:</b> <ul style="list-style-type: none"> <li>Infused by a health care professional</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>50 mg/4 mL single dose vial</li> </ul> | <ul style="list-style-type: none"> <li>Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>Do not freeze</li> <li>Do not shake</li> <li>Store in original carton to protect from light</li> </ul>            | Upper respiratory tract infection, alanine aminotransferase increased, viral infection, aspartate aminotransferase increased, neutrophil count decreased, bronchitis, hypertension and rash |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                             | Manufacturer          | FDA-Approved Indication(s)   | Route of Administration   | Supply Method  | Storage   | Select Common Adverse Reactions  |
|---------------------------------------|-----------------------|--|---|--|---|--|
| <b>Skyrizi</b><br>(risankizumab-rzaa) | AbbVie                | <b>IV/OBI:</b> <ul style="list-style-type: none"> <li>• CD</li> <li>• UC</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>• PsA</li> <li>• PsO</li> </ul> | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Infused by a health care professional</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 600 mg/10 mL single dose vial</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>• 360 mg/2.4 mL OBI</li> <li>• 150 mg/mL PEN</li> <li>• 150 mg/mL PFS</li> </ul> | <ul style="list-style-type: none"> <li>• Store in a refrigerator at 36°F to 46° F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Do not shake</li> <li>• Store in the original cartons to protect from light</li> </ul> | <b>PsO and PsA:</b> upper respiratory infections, headache, fatigue, injection site reactions and tinea infections<br><br><b>CD:</b><br><b>Induction:</b> upper respiratory infections, headache, and arthralgia.<br><b>Maintenance:</b> arthralgia, abdominal pain, injection site reactions, anemia, pyrexia, back pain, arthropathy and urinary tract infection |
| <b>Sotyktu</b><br>(deucravacitinib)   | Bristol Meyers Squibb | <ul style="list-style-type: none"> <li>• PsO</li> </ul>  | Oral  | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 6 mg tablets</li> </ul>  | Store at 68°F to 77°F (20°C to 25°C)  | Upper respiratory infections, blood creatine phosphokinase increased, herpes simplex, mouth ulcers, folliculitis and acne  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                       | Manufacturer | FDA-Approved Indication(s)  | Route of Administration  | Supply Method  | Storage  | Select Common Adverse Reactions   |
|---------------------------------|--------------|---|--|--|--|---|
| <b>Stelara</b><br>(ustekinumab) | Janssen      | <ul style="list-style-type: none"> <li>• CD</li> <li>• PsA (6+)</li> <li>• PsO (6+)</li> <li>• UC</li> </ul>  | <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul><br><b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>IV:</b> <ul style="list-style-type: none"> <li>• 130 mg/26 mL single dose vial</li> </ul><br><b>SC:</b> <ul style="list-style-type: none"> <li>• 45 mg/0.5 mL PFS</li> <li>• 90 mg/mL PFS</li> <li>• 45 mg/0.5 mL single-dose vial</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Do not shake</li> <li>• Store in original carton to protect from light</li> </ul> | <b>PsO:</b> nasopharyngitis, upper respiratory tract infection, headache and fatigue<br><b>CD:</b> Induction: vomiting.<br><b>Maintenance:</b> nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic infection, bronchitis, pruritus, urinary tract<br><b>UC:</b> Induction: nasopharyngitis.<br><b>Maintenance:</b> nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea |
| <b>Taltz</b><br>(ixekizumab)    | Eli Lilly    | <ul style="list-style-type: none"> <li>• AS</li> <li>• Nr-axSpA</li> <li>• PsA</li> <li>• PsO (6+)</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>   | <b>SC:</b> <ul style="list-style-type: none"> <li>• 80 mg/mL PEN</li> <li>• 80 mg/mL PFS</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> <li>• Do not shake</li> </ul>                             | Injection site reactions, upper respiratory tract infections, nausea and tinea infections   |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                           | Manufacturer   | FDA-Approved Indication(s)  | Route of Administration   | Supply Method   | Storage  | Select Common Adverse Reactions   |
|-------------------------------------|----------------|---|---|---|--|---|
| <b>Tremfya</b><br>(guselkumab)      | Janssen        | <ul style="list-style-type: none"> <li>• PsA</li> <li>• PsO</li> </ul>  | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul>  | <b>SC:</b> <ul style="list-style-type: none"> <li>• 100 mg/mL PFS</li> <li>• 100 mg/mL PEN</li> </ul>   | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Store in original carton to protect from light</li> <li>• Do not freeze</li> <li>• Do not shake</li> </ul>                | Upper respiratory infections, headache, injection site reactions, arthralgia, bronchitis, diarrhea, gastroenteritis, tinea infections and herpes simplex infections   |
| <b>Truxima</b><br>(rituximab-abbs)  | Celltrion/Teva | <ul style="list-style-type: none"> <li>• RA</li> </ul>  | <b>IV infusion only:</b> <ul style="list-style-type: none"> <li>• Administer only by a health care professional</li> </ul>  | <b>IV:</b> <ul style="list-style-type: none"> <li>• 100 mg/10 mL single dose vial</li> <li>• 500 mg/50 mL single dose vial</li> </ul>   | <ul style="list-style-type: none"> <li>• Refrigerate vials at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze or shake</li> <li>• Protect from direct sunlight</li> </ul>   | Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events) |
| <b>Tyenne</b><br>(tocilizumab-aazg) | Fresenius Kabi | <ul style="list-style-type: none"> <li>• GCA</li> <li>• PJIA (2+)</li> <li>• RA</li> <li>• SJIA (2+)</li> </ul> | <b>SC Injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> <b>IV infusion:</b> <ul style="list-style-type: none"> <li>• Prepared and infused by a health care professional</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 162 mg/0.9 mL PFS</li> <li>• 162 mg/0.9 mL PEN</li> </ul> <b>IV:</b> <ul style="list-style-type: none"> <li>• 80 mg/4 mL</li> <li>• 200 mg/10 mL</li> <li>• 400 mg/20 mL</li> </ul> | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Store in original packages</li> <li>• Protect from light</li> <li>• Keep syringes dry</li> </ul> | Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions  |

# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                                  | Manufacturer | FDA-Approved Indication(s)   | Route of Administration  | Supply Method  | Storage   | Select Common Adverse Reactions  |
|--|--------------|--|--|--|---|--|
| <b>Velsipity</b><br>(etrasimod)            | Pfizer       | <ul style="list-style-type: none"> <li>• UC</li> </ul>   | Oral   | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 2 mg tablet</li> </ul>   | <ul style="list-style-type: none"> <li>• Store at room temperature at 68°F to 77°F (20°C to 25°C)</li> </ul>  | Headache, elevated liver tests and dizziness   |
| <b>Xeljanz/Xeljanz XR</b><br>(tofacitinib) | Pfizer       | <ul style="list-style-type: none"> <li>• AS</li> <li>• PJIA (2+)</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> </ul>   | Oral   | <b>Oral:</b> <ul style="list-style-type: none"> <li>• Tablets:               <ul style="list-style-type: none"> <li>– 5 mg</li> <li>– 10 mg</li> </ul> </li> <li>• XR tablets:               <ul style="list-style-type: none"> <li>– 11 mg</li> <li>– 22 mg</li> </ul> </li> <li>• Oral Solution:               <ul style="list-style-type: none"> <li>– 1 mg/mL</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>• Store at room temperature at 68°F to 77°F (20°C to 25°C)</li> <li>• Do not repackage</li> </ul>            | RA, PsA, AS and PJIA: upper respiratory tract infection, nasopharyngitis, diarrhea and headache<br><br>UC: nasopharyngitis, elevated cholesterol levels, headache, upper respiratory tract infection, increased blood creatine phosphokinase, rash, diarrhea and herpes zoster |
| <b>Yuflyma</b><br>(adalimumab-aaty)        | Celltrion    | <ul style="list-style-type: none"> <li>• AS</li> <li>• CD (6+)</li> <li>• HS</li> <li>• PJIA (2+)</li> <li>• PsO</li> <li>• PsA</li> <li>• RA</li> <li>• UC</li> <li>• UV</li> </ul> | <b>SC injection:</b> <ul style="list-style-type: none"> <li>• Self-administered</li> </ul> | <b>SC:</b> <ul style="list-style-type: none"> <li>• 40 mg/0.4 mL PFS</li> <li>• 40 mg/0.4 mL PEN</li> <li>• 80 mg/0.8 mL PEN</li> </ul>  | <ul style="list-style-type: none"> <li>• Refrigerate at 36°F to 46°F (2°C to 8°C)</li> <li>• Do not freeze</li> <li>• Protect from light</li> </ul> | Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash   |



# Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

| Drug name                    | Manufacturer         | FDA-Approved Indication(s) | Route of Administration | Supply Method  | Storage  | Select Common Adverse Reactions                                |
|------------------------------|----------------------|----------------------------|-------------------------|--|--|--|
| <b>Zeposia</b><br>(ozanimod) | Bristol Myers Squibb | • UC                       | Oral                    | <b>Oral:</b> <ul style="list-style-type: none"> <li>• 7 Day starter pack: <ul style="list-style-type: none"> <li>– 0.23 mg capsules</li> <li>– 0.46 mg capsules</li> </ul> </li> <li>• 0.92 mg capsules</li> </ul> | <ul style="list-style-type: none"> <li>• Store at 68°F to 77°F (20°C to 25°C)</li> </ul> | Liver test increased, upper respiratory infection and headache |



Visit [cvs.co/ai-list](https://cvs.co/ai-list) or scan the QR code to view electronically. For easy access, you can bookmark this page.

FDA (U.S. Food and Drug Administration). SC (Subcutaneous). IV (Intravenous). PFS (Prefilled syringe). OBI (On-body injector).

AA (Alopecia areata). AD (Atopic dermatitis). aGVHD (Acute graft versus host disease). AOSD (Adult-onset Still's Disease). AS (Ankylosing spondylitis). CAPS (Cryopyrin-associated periodic syndrome). CRS (Cytokine release syndrome). CRSwNP (Chronic rhinosinusitis with nasal polyposis). DIRA (Deficiency of IL-1-receptor antagonist). EoE (Eosinophilic esophagitis). ERA (Enthesitis-related arthritis). FCAS (Familial cold autoinflammatory syndrome). FMF (Familial Mediterranean fever). GCA (Giant cell arteritis). HIDS/MKD (Hyperimmunoglobulin D syndrome / mevalonate kinase deficiency). HS (Hidradenitis Suppurativa). JIA (Juvenile idiopathic arthritis). LN (Lupus Nephritis). MWS (Muckle-Wells syndrome). NOMID (Neonatal-onset multisystem inflammatory disease). Nr-axSpA (Non-radiographic axial spondyloarthritis). PJIA (Polyarticular Juvenile idiopathic arthritis). PMR (Polymyalgia rheumatica). PN (Prurigo nodularis). PsA (Psoriatic arthritis). PsO (Plaque psoriasis). RA (Rheumatoid arthritis). RP (Recurrent pericarditis). SJIA (Systemic Juvenile idiopathic arthritis). SLE (Systemic lupus erythematosus). SSc-ILD (Systemic sclerosis - Interstitial lung disease). TRAPS (Tumor necrosis factor receptor-associated periodic syndrome). UC (Ulcerative colitis). UV (Uveitis).

The information contained in this communication is provided in summary form. It is not intended for use as the sole basis of clinical treatment, as a substitute for reading the original research, nor as a substitute for the knowledge, skill and judgment of the medical provider. This list is subject to change. CVS Specialty assumes no liability whatsoever for the information provided or for any diagnosis or treatment made as a result of this information. The information noted in this brochure was taken from the manufacturers' prescribing information. It does not contain all information needed to use these products safely and effectively. See the manufacturers' package insert for full prescribing information. CVS Specialty does not provide medical advice and does not recommend a particular treatment or brand. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Specialty.